Synthesis and Coordination Chemistry of a Novel Phosphinimine Phosphine

Chung-Yuan Liu, Der-Yi Chen, Ming-Chu Cheng, Shie-Ming Peng, and Shiuh-Tzung Liu*

Department of Chemistry, National Taiwan University, Taipei, Taiwan 106, Republic of China

Received November 16, 1994[®]

The incorporation of a phosphinimino moiety into phosphine ligands to form a new bidentate $\text{Ph}_3\text{P}=\text{N}(\text{CH}_2)_3\text{PPh}_2$ (**1**) is reported. Compound **1** can act as a σ -N and σ -P donor ligand or react with metal carbonyls to form isocyanide complexes. Complexes of $(\text{CO})_4\text{M}-(1\text{-P,N})$ [$\text{M} = \text{Mo}$ (**2**), W (**3**)] were prepared from the reaction of $\text{Et}_4\text{N}[\text{M}(\text{CO})_5\text{Br}]$ with **1** in refluxing THF solution, whereas the isocyanide complexes of $(\text{CO})_5\text{MCN}(\text{CH}_2)_3\text{PPh}_2$ [$\text{M} = \text{Cr}$ (**4**), Mo (**5**), W (**6**)] are obtained from the reaction of $\text{M}(\text{CO})_6$ with **1** at 25 °C. The deoxygenation nature is also shown in the reaction of $\text{CpFe}(\text{CO})_2\text{I}$, $\text{CpRu}(\text{CO})_2\text{I}$, $\text{Re}(\text{CO})_5\text{Br}$, and $\text{Re}_2(\text{CO})_{10}$ with **1** to form the corresponding isocyanide complexes $[\text{CpFe}(\text{CO})\{\text{CN}(\text{CH}_2)_3\text{PPh}_2\text{-C,P}\}]\text{I}$ (**7**), $\text{CpRu}(\text{CO})\{\text{CN}(\text{CH}_2)_3\text{PPh}_2\text{-C}\}$ (**10**), $\text{BrRe}(\text{CO})_{5-n}\{\text{CN}(\text{CH}_2)_3\text{PPh}_2\text{-C}\}_n$ [$n = 1$ (**12**), 2 (**14**)], and $\text{Re}_2(\text{CO})_9\{\text{CN}(\text{CH}_2)_3\text{PPh}_2\text{-C}\}$ (**15**), respectively. The free phosphine of **10** underwent exchange with one triphenylphosphine ligand in $\text{CpRu}(\text{PPh}_3)_2\text{Cl}$ to yield the binuclear species $[\text{CpRu}(\text{PPh}_3)\text{Cl}\{\text{P}(\text{Ph}_2)(\text{CH}_2)_3\text{NC}\}\text{Ru}(\text{CO})\text{CpI}]$ (**11**), whereas intramolecular ligand substitution occurred in both **12** and **15** to give $\text{Br}(\text{CO})_3\text{Re}\{\text{CN}(\text{CH}_2)_3\text{PPh}_2\text{-C,P}\}$ (**13**) and $\{\mu\text{-CN}(\text{CH}_2)_3\text{PPh}_2\}\text{Re}_2(\text{CO})_8$ (**16**). Reaction of $(\text{COD})\text{PdCl}_2$ with **1** produced the complex $(1\text{-P,N})\text{PdCl}_2$ (**18**), in which **1** acts as a σ -N, σ -P bidentate ligand. X-ray crystal structural analysis of $[\text{CpFe}(\text{CO})\{\text{CN}(\text{CH}_2)_3\text{PPh}_2\text{-C,P}\}]\text{PF}_6$ (**7a**) and **18** confirmed the formulation of both complexes. The $\text{C}\equiv\text{N}$ bond distance [1.20(3) Å] of **7a** is greater than those in the related iron–isocyanide complexes; the angle $\text{C}-\text{N}-\text{C}$ [141(2)°] deviates from 180°, indicating that the resonance contribution of $\text{Fe}=\text{C}=\text{N}-$ is more important than that of $\text{Fe}-\text{C}\equiv\text{N}-$. These observations are consistent with spectral data, the smaller infrared stretching wavenumber (2089 cm^{-1}), and the greater shift (^{13}C NMR δ 183.7 ppm) of the isocyanide moiety of **7**.

Introduction

The nature of the highly polar P–N bond in phosphinimine makes a ligand of this kind versatile in both coordination and organometallic chemistry.^{1–14} The nitrogen atom of phosphinimine is able to act as a two-

electron donor for complexation with various metal ions^{2,3} and as a four-electron donor (bridging mode) in $\text{Mo}_2(\text{CO})_6(\text{Ph}_3\text{PNH})_3$.⁴ Furthermore, phosphinimines can react with metal carbonyl complexes⁶ and metal oxides⁷ to produce isocyanide and nitrene functions, respectively, by elimination of phosphine oxide.

Incorporation of a phosphinimino moiety with other donor atoms to form a chelate bidentate has been reported for $9,10\text{-H}_2\text{NC}_{14}\text{H}_8\text{N}=\text{PPh}_3$,⁸ $o\text{-(HO)C}_6\text{H}_4\text{N}=\text{PPh}_3$,⁹ $8\text{-(Ph}_3\text{P}=\text{N-})\text{C}_9\text{H}_6\text{N}$,¹⁰ $(\eta^1\text{-C}_5\text{Me}_5)\text{P}=\text{N}(t\text{-Bu})$,¹¹ $\text{Ph}_3\text{P}=\text{NC}_6\text{H}_4\text{N}=\text{PPh}_3$,^{7b,12} $\text{Ph}_2\text{P}(\text{=NR})\text{CH}_2\text{PPh}_2(\text{=NR})$,¹³ and $\text{PhN}=\text{P}(\text{Ph}_2)\text{CH}=\text{CH}(p\text{-Tol})(\text{NH}_2)$.¹⁴ Among these bidentate ligands, only one phosphine–phosphinimine, $\text{Ph}_2\text{PCH}_2\text{PPh}_2(\text{=NSiMe}_3)$, is reported,¹⁵ for which the coordination mode toward metal ions is confined due to the short carbon chain. We recently showed that the phosphinimine function is stable toward the phosphide nucleophile, which allows us to prepare the potential bidentate phosphino–phosphinimine ligands **1**.¹⁶ Analogous to $\text{Ph}_2\text{PCH}_2\text{PPh}_2(\text{=NSiMe}_3)$, one expects compound **1** to be able to coordinate in a σ -N and σ -P chelate mode toward metal ions and that compound **1** would react

* Abstract published in *Advance ACS Abstracts*, March 15, 1995.

(1) Dehnicke, K.; Strahle, J. *Polyhedron* **1989**, *8*, 707.
(2) Choukroun, R.; Gervais, D.; Dilworth, J. R. *Transition Met. Chem.* **1979**, *4*, 249.

(3) (a) Sleiman, H. F.; Mercer, S.; McElwee-White, L. *J. Am. Chem. Soc.* **1989**, *111*, 8007. (b) Briggs, E. M.; Brown, G. W.; Jiricny, J. *J. Inorg. Nucl. Chem.* **1979**, *41*, 667.

(4) Miller, J. S.; Visscher, M. O.; Caulton, K. G. *Inorg. Chem.* **1974**, *13*, 1632.

(5) Roesky, H. W.; Munzenberg, J.; Bohra, R.; Noltemeyer, M. *J. Organomet. Chem.* **1991**, *418*, 339.

(6) (a) Alper, H.; Partis, R. A. *J. Organomet. Chem.* **1972**, *35*, C40. (b) Mirkin, C. A.; Lu, K.-L.; Geoffroy, G. L. *J. Am. Chem. Soc.* **1989**, *111*, 7279. (c) Lin, Y.-W.; Gau, H.-M.; Wen, Y.-S.; Lu, K.-L. *Organometallics* **1992**, *11*, 1445.

(7) (a) Maatta, E. A.; Haymore, B. L.; Wentworth, R. A. *D. Inorg. Chem.* **1980**, *19*, 1055. (b) Maatta, E. A.; Kim, C. *Inorg. Chem.* **1989**, *28*, 623. (c) Chatt, J.; Rowe, G. A. *J. Chem. Soc. A* **1969**, 2288. (d) Chong, A. O.; Oshima, K.; Sharpless, K. B. *J. Am. Chem. Soc.* **1977**, *99*, 3420.

(8) Dapporto, P.; Denti, G.; Dolcetti, G.; Ghedini, M. *J. Chem. Soc., Dalton Trans.* **1983**, 779.

(9) Charalambous, J.; Kensett, M. J.; Jenkins, J. M. *J. Chem. Res., Synop.* **1982**, 306.

(10) Saravanamuthu, A.; Ho, D. M.; Kerr, M. E.; Fitzgerald, C.; Bruce, M. R. M.; Bruce, A. E. *Inorg. Chem.* **1993**, *32*, 2202.

(11) Gudat, D.; Niecke, E.; Krebs, B.; Dartmann, M. *Organometallics* **1986**, *5*, 2376.

(12) Herring, D. L. *J. Org. Chem.* **1961**, *26*, 3998.

(13) Imhoff, P.; van Asselt, R.; Elsevier, C. J.; Zoutberg, M. C.; Stam, C. H. *Inorg. Chim. Acta* **1991**, *184*, 73.

(14) Fernandez, M. J.; Del Val, J. J.; Oro, L. A.; Palacios, F.; Barluenga, J. *Polyhedron* **1987**, *6*, 1999.

(15) (a) Katti, K. V.; Cavell, R. G. *Organometallics* **1988**, *7*, 2236. (b) Katti, K. V.; Cavell, R. G. *Inorg. Chem.* **1989**, *28*, 413. (c) Katti, K. V.; Cavell, R. G. *Inorg. Chem.* **1989**, *28*, 3033. (d) Katti, K. V.; Cavell, R. G. *Organometallics* **1989**, *8*, 2147. (e) Katti, K. V.; Batchelor, R. J.; Einstein, F. W. B.; Cavell, R. G. *Inorg. Chem.* **1990**, *29*, 808. (f) Katti, K. V.; Cavell, R. G. *Organometallics* **1991**, *10*, 539.

(16) Liu, S.-T.; Liu, C.-Y. *J. Org. Chem.* **1992**, *57*, 6079.

Figure 1. ORTEP view of the cation of complex **7a** showing 50% probability thermal ellipsoids. Two independent molecules exist in the unit cell.

Scheme 1

with a metal–carbonyl ligand to form a new ligand $\text{CN}-(\text{CH}_2)_3\text{PPh}_2$ of bidentate type in either a bridging or a chelating mode. Here we report the results of coordination chemistry of **1** toward various metal complexes.

Results and Discussion

The ligand $\text{Ph}_3\text{P}=\text{N}-(\text{CH}_2)_3\text{PPh}_2$ was obtained after isolation of the intermediate in the synthesis of $\text{H}_2\text{N}-(\text{CH}_2)_3\text{PPh}_2$ (Scheme 1).¹⁶ This phosphino–phosphinimine **1** is stable in air but sensitive to moisture. The ^{31}P NMR spectrum of the new ligand displays two singlets at -16.0 and 4.3 ppm in benzene- d_6 for diphenylphosphino and iminophosphorane moieties, respectively. The ^{31}P chemical shift corresponding to the moiety of $\text{Ph}_3\text{P}=\text{N}-$ is varied from nonpolar solvent to polar solvent. Thus the chemical shift of the phosphinimino moiety is 12.1 ppm in CDCl_3 , indicating the polar nature of the P–N bond.

Addition of **1** to a stirred tetrahydrofuran solution of $\text{NEt}_4[\text{M}(\text{CO})_5\text{Br}]$ ($\text{M} = \text{Mo}, \text{W}$) at refluxing temperature results in the formation of chelate complexes $(\text{CO})_4\text{M}-(1-P,N)$ [**2**, $\text{M} = \text{Mo}$; **3**, $\text{M} = \text{W}$]. Both complexes were isolated as solids stable in air and characterized by spectral methods. The carbonyl stretching wavenumbers of the complexes occur in the 2010 – 1900 cm^{-1} region, which is in accord with a tetracarbonyl species. The ^{31}P NMR spectrum displays two sets of doublets [39.4 (d, $J = 14$ Hz), 16.3 (d, $J = 14$ Hz) ppm for **2** and 37.9 (d, $J = 18$ Hz), 2.67 (dt, $J_{\text{P-P}} = 18$ Hz, $J_{\text{P-W}} = 224$ Hz) ppm for **3**] due to phosphorus atoms of two kinds coupled to each other. Both the coordination chemical shifts and long range phosphorus–phosphorus couplings verify the existence of the $\text{Ph}_3\text{P}=\text{N}-$ moiety in the complexes. The infrared spectra of both **2** and **3** exhibit

Table 1. Infrared Absorptions and ^{13}C NMR Chemical Shifts of Isocyanide Complexes^a

complex	$\nu_{\text{CN}}, \text{cm}^{-1}$	^{13}C NMR, ppm	ref
4	2173	162.5	this work
5	2173	153.4	this work
6	2175	143.0	this work
7	2089	183.7 (d, $J_{\text{P-C}} = 35$ Hz)	this work
$[\text{CpFe}(\text{PPh}_3)(\text{CO})(\text{CNCH}_3)]\text{BF}_4$	2194		20
10	2170	145.1	this work
11	2170	144.9, 144.8 ^c	this work
12	2222	129.2	this work
$\text{ReBr}(\text{CO})_4(\text{CNPr})$	2221	129.4	21
13	2138	163 (d, $J_{\text{P-C}} = 6.8$ Hz)	this work
<i>fac</i> - $\text{Re}(\text{CO})_3(\text{CN}^i\text{Bu})(\text{PPh}_2\text{Me})\text{Br}$	2186		23
<i>fac</i> - $\text{Re}(\text{CO})_3(\text{CN}^i\text{Bu})(\text{PPh}_3)\text{Br}$	2181		23
14	2219, 2192	134.3	this work
$\text{ReBr}(\text{CO})_3(\text{CNPr})_2$	2219, 2197	134.5	21
15	2187	134.5	this work
$\text{Re}_2(\text{CO})_9(\text{CN}^i\text{Bu})$	2173 ^d		22
16	2156 ^b	178.9	this work
17	2221	130.7	this work

^a Recorded in CH_2Cl_2 for IR and in CDCl_3 for ^{13}C NMR, unless otherwise noted. ^b KBr. ^c Diastereomers. ^d In hexane.

no band that could be assigned to $\text{C}\equiv\text{N}-$ modes, in agreement with the structural formulation. Apparently, compound **1** acts as a simple chelate ligand (a $\sigma\text{-N}$ and $\sigma\text{-P}$ chelate mode) in $\text{Mo}(0)$ and $\text{W}(0)$ complexes.

Deoxygenation of the carbonyl group took place in the reaction of **1** with $\text{M}(\text{CO})_6$ at room temperature to yield the corresponding isocyanide complex $(\text{CO})_5\text{M}[\text{CN}(\text{CH}_2)_3\text{PPh}_2\text{-C}]$ [**4**, $\text{M} = \text{Cr}$; **5**, $\text{M} = \text{Mo}$; **6**, $\text{M} = \text{W}$]. The structural determination of **4**–**6** was confirmed by infrared and NMR spectra. Infrared spectra of these series complexes exhibit characteristic absorptions at ca. 2174 cm^{-1} for the coordinated isocyanide group and

Table 2. Atomic Parameters and Thermal Parameters of 7a

	x	y	z	B _{eq} , Å ²
Fe	0.8039(4)	0.10504(21)	0.3151(3)	4.74(20)
FeA	-0.2101(4)	0.60450(21)	0.1790(3)	4.93(21)
P	0.9299(8)	0.0482(4)	0.2060(5)	4.5(4)
PA	-0.0815(8)	0.5491(4)	0.2933(5)	4.5(4)
C1	0.693(3)	0.0222(15)	0.3925(16)	5.8(15)
C2	0.644(3)	0.0930(17)	0.4132(22)	9.4(22)
C3	0.764(3)	0.1324(15)	0.4491(18)	6.3(16)
C4	0.891(3)	0.0897(16)	0.4446(17)	6.7(16)
C5	0.8376(25)	0.0201(14)	0.4105(15)	5.0(14)
C6	1.000(3)	0.1108(13)	0.1170(16)	6.4(17)
C7	1.138(3)	0.1578(13)	0.1465(18)	7.1(17)
C8	1.106(3)	0.2374(15)	0.1836(18)	6.6(18)
N	0.9989(22)	0.2295(10)	0.2562(13)	5.6(12)
C9	0.9133(25)	0.1844(14)	0.2854(18)	5.6(15)
C10	0.668(3)	0.1222(15)	0.2373(23)	9.4(21)
O10	0.5616(20)	0.1276(12)	0.1930(13)	9.1(13)
C11	0.818(3)	-0.0158(14)	0.1395(15)	4.9(6)
C12	0.778(3)	-0.0877(15)	0.1793(16)	5.7(7)
C13	0.678(3)	-0.1348(15)	0.1320(16)	5.9(7)
C14	0.622(3)	-0.1131(15)	0.0470(16)	5.9(7)
C15	0.659(3)	-0.0480(16)	0.0074(18)	6.9(8)
C16	0.753(3)	0.0006(15)	0.0554(17)	6.4(7)
C21	1.0888(24)	-0.0072(13)	0.2416(14)	4.1(6)
C22	1.176(3)	0.0137(14)	0.3150(16)	5.5(7)
C23	1.301(3)	-0.0266(15)	0.3377(17)	6.5(7)
C24	1.346(3)	-0.0803(15)	0.2771(16)	5.9(7)
C25	1.265(3)	-0.1006(14)	0.2047(15)	5.3(7)
C26	1.1355(24)	-0.0681(13)	0.1824(14)	3.9(6)
C1A	-0.109(3)	0.5891(15)	0.0512(17)	6.6(16)
C2A	-0.246(4)	0.6364(15)	0.0397(19)	8.5(20)
C3A	-0.364(3)	0.5954(15)	0.0664(19)	7.5(17)
C4A	-0.307(3)	0.5193(15)	0.0988(16)	6.5(16)
C5A	-0.162(3)	0.5223(14)	0.0827(14)	4.5(14)
C6A	-0.009(3)	0.6139(14)	0.3840(16)	5.3(15)
C7A	0.124(3)	0.6643(14)	0.3626(18)	6.1(16)
C8A	0.074(3)	0.7477(14)	0.3228(19)	6.9(18)
NA	-0.029(3)	0.7323(11)	0.2462(13)	7.1(14)
C9A	-0.104(3)	0.6870(14)	0.2157(16)	6.3(16)
C10A	-0.350(3)	0.6229(14)	0.2482(17)	5.2(14)
O10A	-0.4500(20)	0.6329(12)	0.2949(12)	8.7(13)
C11A	-0.1901(25)	0.4821(13)	0.3532(15)	4.6(6)
C12A	-0.252(3)	0.4972(15)	0.4341(16)	5.6(7)
C13A	-0.344(3)	0.4453(17)	0.4784(19)	8.1(8)
C14A	-0.369(3)	0.3783(16)	0.4340(18)	7.1(8)
C15A	-0.319(3)	0.3588(16)	0.3554(18)	7.6(8)
C16A	-0.222(3)	0.4086(14)	0.3099(16)	5.6(7)
C21A	0.0803(24)	0.4988(13)	0.2652(14)	3.9(6)
C22A	0.129(3)	0.4378(15)	0.3219(16)	5.7(7)
C23A	0.268(3)	0.4053(15)	0.3038(16)	5.8(7)
C24A	0.344(3)	0.4237(16)	0.2312(17)	6.6(7)
C25A	0.297(3)	0.4861(16)	0.1718(17)	6.7(7)
C26A	0.165(3)	0.5195(15)	0.1943(16)	5.5(7)
P1	0.2762(10)	0.7122(5)	-0.0048(6)	7.3(5)
F1	0.373(3)	0.7305(10)	0.0772(15)	15.7(16)
F2	0.189(3)	0.6917(14)	-0.0872(16)	19.2(20)
F3	0.2680(23)	0.6285(9)	0.0263(13)	12.4(14)
F4	0.2832(24)	0.7966(12)	-0.0284(18)	18.1(20)
F5	0.1334(22)	0.7286(12)	0.0395(17)	15.5(17)
F6	0.4217(23)	0.6960(13)	-0.0516(14)	15.4(16)
P2	0.6984(10)	0.7900(5)	0.4896(6)	7.0(5)
F7	0.643(3)	0.7798(10)	0.5868(12)	13.4(15)
F8	0.752(3)	0.8018(12)	0.3930(12)	15.2(16)
F9	0.703(3)	0.8747(9)	0.5146(15)	14.8(16)
F10	0.6975(23)	0.7042(10)	0.4713(15)	13.5(15)
F11	0.8552(20)	0.7828(12)	0.5213(15)	14.3(16)
F12	0.5378(21)	0.7955(12)	0.4558(13)	13.4(14)

Table 3. Selected Bond Distances (Å) and Bond Angles (deg)

Complex 7a			
Fe-P	2.229(9)	FeA-PA	2.229(9)
Fe-C9	1.77(2)	FeA-C9A	1.80(2)
Fe-C10	1.67(3)	FeA-C10A	1.69(2)
C9-N	1.20(3)	C9A-NA	1.12(3)
N-C8	1.47(3)	NA-C8A	1.45(3)
C10-O10	1.16(3)	C10A-O10A	1.17(3)
C10-Fe-P	90(1)	C10A-FeA-PA	92.2(8)
C10-Fe-C9	96(1)	C10A-FeA-C9A	95(1)
C9-Fe-P	82.0(8)	C9A-FeA-PA	82.7(8)
Fe-C9-N	169(2)	FeA-C9A-NA	171(2)
C9-N-C8	141(2)	C9A-NA-C8A	142(2)
Fe-C10-O10	169(3)	FeA-C10A-O10A	177(2)
Complex 18			
Pd-Cl1	2.371(2)	Pd-N	2.067(5)
Pd-Cl2	2.277(2)	P2-N	1.575(6)
Pd-P1	2.203(2)		
Cl1-Pd-Cl2	91.17(8)	P1-Pd-N	83.5(2)
Cl2-Pd-P1	92.19(8)	P2-N-C8	119.4(4)
Cl2-Pd-N	175.1(2)	P2-N-Pd	125.6(3)
Cl1-Pd-P1	171.17(8)	C8-N-Pd	112.8(4)
Cl1-Pd-N	93.4(2)		

oxide.⁶ It appears that the carbonyl ligand of M(CO)₆ is more reactive toward nucleophilic attack by the iminophosphorane than that of [M(CO)₅Br]⁻. Such a difference is believed due to the poorer M-C back-bonding of M(CO)₆ compared with that of [M(CO)₅Br]⁻.

Reaction of phosphine-phosphinimine **1** with CpFe(CO)₂I [Cp = η⁵-C₅H₅] in toluene at 25 °C yielded a yellow precipitate {CpFe(CO)[CNCH₂CH₂CH₂PPh₂-P,C]}I (**7**) and triphenylphosphine oxide; the desired complex was purified by extraction of Ph₃P=O with toluene from the reaction mixture. Anionic metathesis of **7** with NH₄PF₆ gave {CpFe(CO)[CNCH₂CH₂CH₂-PPh₂-P,C]}PF₆ (**7a**) as a crystalline solid upon recrystallization. The phosphinimine function of **1** acts as a deoxygenating agent, which reacts with the carbonyl ligand to form an isocyanide moiety. In the infrared spectrum of **7**, the stretching wavenumbers corresponding to isocyanide (2089 cm⁻¹) and carbonyl (1998 cm⁻¹) were observed, clearly indicating the existence of both moieties around the metal center. An ORTEP plot of the cation of **7a** is illustrated in Figure 1. Atomic coordinates and selected bond distances and angles are listed in Tables 2 and 3, respectively. The iron atom displays a slightly distorted octahedral geometry with one face occupied by the cyclopentadienyl moiety. The Fe-C9 distance [1.77(2) Å] is smaller and the C9-N distance [1.20(3) Å] is larger than the average bond distance of Fe-C (1.84 Å) and C-N (1.17 Å) in complexes of Fe(*t*-BuNC)₅,¹⁷ Fe(C₆H₅)(CO)₂(EtNC),¹⁸ and Fe₂(EtNC)₉,¹⁹ respectively. The angle C9-N-C8 [141-(2)°] is much deviated from 180°. All this information indicates that the nitrogen atom is no longer restrained in a linear environment, which implies that the reso-

at ca. 2070 and 1955 cm⁻¹ for the pentacarbonylmetal moiety. The ¹³C NMR shifts of the isocyanide carbon appear at 162.5 ppm for **4**, 153.4 ppm for **5**, and 143.0 ppm for **6**, respectively. Both IR and NMR data indicate the existence of an isocyanide function. The formation of isocyanide complex is presumably proceeded by the nucleophilic attack of iminophosphorane at the carbonyl ligand followed by the elimination of triphenylphosphine

(17) Bassett, J.-M.; Berry, D. E.; Barker, G. K.; Green, M.; Howard, J. A. K.; Stone, F. G. A. *J. Chem. Soc., Dalton Trans.* **1979**, 1003.

(18) Yamamoto, Y. *Coord. Chem. Rev.* **1980**, 32, 193.

(19) Bassett, J.-M.; Barker, G. K.; Green, M.; Howard, J. A. K.; Stone, F. G. A.; Wolsey, W. C. *J. Chem. Soc., Dalton Trans.* **1981**, 219.

(20) Angelici, R. J.; Christian, P. A.; Dombek, B. D.; Pfeiffer, G. A. *J. Organomet. Chem.* **1974**, 67, 287.

(21) Chen, L.-C.; Chen, M.-Y.; Chen, J.-H.; Wen, Y.-S.; Lu, K.-L. *J. Organomet. Chem.* **1992**, 425, 99.

(22) Harris, G. W.; Coville, N. J. *Organometallics* **1985**, 4, 908.

(23) Leins, A. E.; Coville, N. J. *J. Organomet. Chem.* **1994**, 464, 183.

nance form $\text{Fe}=\text{C}=\text{N}-$ in the coordinated isocyanide group is much more important than $\text{Fe}-\text{C}\equiv\text{N}-$. This observation is also consistent with the smaller stretching wavenumber of isocyanide (2089 cm^{-1}) in **7** (Table 1) compared with $[\text{CpFe}(\text{PPh}_3)(\text{CO})(\text{CNCH}_3)]\text{BF}_4$ (2194 cm^{-1}). Complex **7** readily reacted with propylamine to give carbene complex **8**. The structure of this carbene metal complex was determined by spectral and elemental analyses (Experimental Section).

The chemistry of **1** toward ruthenium(II) complexes is summarized in Scheme 2. As expected, the ligand substitution reaction of $\text{CpRu}(\text{PPh}_3)_2\text{Cl}$ by the phosphino site of **1** took place immediately to yield the simple substituted complex **9**, but further substitution to form a $\sigma\text{-N}$ and $\sigma\text{-P}$ chelate complex did not occur, even under severe reaction conditions. Reaction of **1** with $\text{CpRu}(\text{CO})_2\text{I}$ produced isocyanide complex **10** accompanied by formation of triphenylphosphine oxide. Unlike the iron complex, the diphenylalkylphosphine moiety in **10** remained as a free donor. Attempts to prepare a chelating mode of $\text{CN}(\text{CH}_2)_3\text{PPh}_2$ complex, like the iron

analog, failed. Nevertheless, the uncoordinating donors in both **9** and **10** become an advantage in preparation of a binuclear species. Thus reaction of either complex **9** with $\text{CpRu}(\text{CO})_2\text{I}$ or complex **10** with $\text{CpRu}(\text{PPh}_3)_2\text{Cl}$ yielded a bridging diruthenium complex **11**. As both ruthenium centers are chiral, the diastereomeric pair of complexes **11** were observed in ^{31}P NMR spectrum: one is 44.1 and 35.7 ppm; the other is 44.0 and 35.8 ppm. All these absorptions appear as doublets with phosphorus-phosphorus coupling constant 41 Hz. Attempts to separate these isomers by chromatography and recrystallization were unsuccessful.

The ruthenium complexes were characterized by spectral and elemental analyses. The ^{31}P NMR spectrum of complex **9** shows three sets of resonances: 44.1 (d, $J = 41\text{ Hz}$, PPh_3); 36.8 (d, $J = 41\text{ Hz}$, $-\text{PPh}_2$), and 13.0 (br, $-\text{N}=\text{PPh}_3$) ppm, in which the broad peak at 13.0 ppm corresponds to the free phosphinimine site. The naturally abundant ratio of ^{35}Cl and ^{37}Cl in the FAB mass spectrum of **9** indicates that the chloride is a coordinated ligand around the metal ion. The ^{31}P NMR spectrum of complex **10** shows only a signal due to a free diphenylalkylphosphine at $\delta -17.3\text{ ppm}$, thus indicating that the phosphine moiety did not coordinate to the metal center and the phosphinimine function disappeared. Both infrared and ^{13}C NMR spectra of **10** show the presence of a coordinated isocyanide (2107 cm^{-1}) and a carbonyl ligand [IR 2107 cm^{-1} (NC) and 1973 cm^{-1} (CO); ^{13}C NMR δ 145.1 ppm (CN) and 199.5 ppm (CO)]. On this basis with elemental analysis the ruthenium complexes **10** and **11** were assigned as formulated.

When $\text{Re}(\text{CO})_5\text{Br}$ reacted with **1** under thermal conditions, deoxygenation at the carbonyl ligand occurred in dichloromethane solution to form an isocyanide complex **12**. The phosphine site of **1** in complex **12** remained a free donor, but it underwent further substitution with carbonyl ligands to give **13**. Treatment of **12** with **1** in a further equimolar quantity provided the diisocyanide

Scheme 3

species **14** that has two free phosphine donors away from the metal center. The structures of **12**–**14** were confirmed by their spectral properties (see Experimental Section). The infrared stretching wavenumber of the isocyanide moiety (2138 cm^{-1}) in **13** is smaller than that of the other $\text{Re}-\text{CN}$ species and the ^{13}C NMR shift of isocyanide carbon (δ 163 ppm) is also greater than those of related species (Table 1), similar to complex **7** due to ring strain.

Reaction of phosphino-phosphinimine **1** with dirhenium decacarbonyl also yielded isocyanide complex **15** in which the phosphino site stays uncoordinated. On irradiation of **15** in chloroform with light, the phosphine exchanged a carbonyl ligand to form a bridging binuclear complex **16** (Scheme 3). Because of the ring strain of **16**, the properties of the coordinated isocyanide ligand in both infrared and ^{13}C NMR spectra show the same trend as in complexes **7** and **13**: a smaller wavenumber of the infrared stretching mode of $\text{CN}-$ and a considerable downfield chemical shift of ^{13}C NMR of $\text{CN}-$. The lack of phosphorus-carbon coupling of the isocyanide carbon supports assignment of a structure in which $\text{CN}(\text{CH}_2)_3\text{PPh}_2$ binds in a bridging mode, not a chelating one. Upon treatment of **16** with bromine, the cleavage of the metal-metal bond of **16** occurred to yield $(\text{CO})_4\text{Re}\{\mu_2-\text{CNCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2-\text{C},\text{P}\}\text{Re}(\text{CO})_4\text{Br}$ (**17**). As the ring strain lessens, the spectral data for the coordinated isocyanide ligand of **17** exhibit the normal features; i.e. both the infrared absorption and ^{13}C NMR chemical shift of the isocyanide group are similar to those of non-chelate complexes (Table 1).

Addition of (cyclooctadiene) PdCl_2 to $\text{Ph}_3\text{P}=\text{N}(\text{CH}_2)_3\text{PPh}_2$ (**1**) in dichloromethane resulted in formation of palladium complex **18** as a yellow solid. This compound is stable, even in the presence of moisture. Complex **18** in chloroform- d_1 solution exhibited a pair of doublets in the ^{31}P NMR spectrum at 17.2 and 40.5 ppm with a coupling constant of 3.3 Hz, clearly indicating that both phosphinimine and phosphine are coordinated at the metal center. The formulation of **18** is further confirmed with X-ray structural analysis.

The results of an X-ray structural analysis of **18** are depicted in Figure 2 and in Tables 3 and 4. The coordination environment about the metal center in **18** is slightly distorted square-planar with the two chloride ligands situated in a *cis* fashion. All bond distances and bond angles in **18** are in the normal range. The $\text{Pd}-\text{Cl}(1)$ bond [$2.371(2)\text{ \AA}$] is notably longer than $\text{Pd}-\text{Cl}(2)$ [$2.277(2)\text{ \AA}$] which is an indication of the *trans* influence of phosphorus versus nitrogen donors. Attempts to prepare nickel and platinum analogs of **18** were unsuccessful.

Summary

We have demonstrated various reactions of the phosphinimine-phosphine $\text{Ph}_3\text{P}=\text{N}(\text{CH}_2)_3\text{PPh}_2$ (**1**) toward metal complexes. For coordination of phosphinimine, compound **1** acts as a σ -P donor (**I**), or in a σ -N and σ -P chelate fashion (**II**) (Chart 1). For deoxygenation of carbonyl, compound **1** becomes an isocyanide-phosphine ligand $\text{CN}(\text{CH}_2)_3\text{PPh}_2$, which acts as a σ -C donor (**III**), a σ -C and σ -P chelate (**IV**), a σ -C and σ -P bridging mode with a metal-metal bond (**V**), and a σ -C and σ -P bridging mode without a metal-metal bond (**VI**).

Experimental Section

General Information. Nuclear magnetic resonance spectra were recorded on either a Bruker AC-E 200 or a AM-300 spectrometer. For ^{31}P NMR spectra, the chemical shifts are given in parts per million (δ) relative to 85% H_3PO_4 . Infrared spectra were measured on a Biorad FT-30 instruments. The

Figure 2. ORTEP diagram of palladium complex 18 showing 50% probability thermal ellipsoids.**Table 4.** Atomic Coordinates and Thermal Parameters of 18

	<i>x</i>	<i>y</i>	<i>z</i>	<i>B</i> _{eq} , Å ²
Pd	0.40595(5)	0.76395(4)	0.05976(3)	2.54(3)
Cl1	0.22543(19)	0.70733(15)	−0.01970(10)	4.61(11)
Cl2	0.52498(20)	0.78297(15)	−0.02029(10)	4.66(11)
P1	0.55445(19)	0.83510(14)	0.13482(10)	2.83(10)
P2	0.26680(19)	0.65156(15)	0.16298(10)	2.98(10)
N	0.3129(5)	0.7467(4)	0.13834(24)	2.7(3)
C6	0.4730(7)	0.9220(5)	0.1750(4)	4.2(5)
C7	0.3599(8)	0.8910(6)	0.2035(5)	6.4(6)
C8	0.2576(7)	0.8336(5)	0.1581(4)	4.1(4)
C12	0.6423(6)	0.7590(5)	0.1982(3)	3.1(4)
C12	0.6773(7)	0.6732(5)	0.1780(4)	3.8(4)
C13	0.7457(8)	0.6116(6)	0.2223(4)	5.9(5)
C14	0.7754(8)	0.6339(7)	0.2878(4)	6.5(6)
C15	0.7430(7)	0.7182(6)	0.3104(4)	5.3(5)
C16	0.6774(6)	0.7817(6)	0.2640(4)	4.1(4)
C21	0.6784(7)	0.9023(5)	0.1059(4)	3.2(4)
C22	0.6451(7)	0.9803(5)	0.0694(4)	4.1(4)
C23	0.7382(8)	1.0339(6)	0.0477(4)	5.0(5)
C24	0.8623(8)	1.0063(6)	0.0621(5)	6.2(6)
C25	0.8989(8)	0.9304(7)	0.0984(5)	7.6(7)
C26	0.8097(7)	0.8747(6)	0.1205(5)	5.5(5)
C31	0.2620(7)	0.6629(5)	0.2483(4)	3.3(4)
C32	0.3749(7)	0.6830(6)	0.2937(4)	4.1(4)
C33	0.3779(8)	0.6959(6)	0.3611(4)	5.6(5)
C34	0.2595(8)	0.6904(6)	0.3828(4)	6.2(6)
C35	0.1490(7)	0.6709(6)	0.3386(4)	5.4(5)
C36	0.1471(7)	0.6569(6)	0.2726(4)	4.2(4)
C41	0.1064(7)	0.6102(5)	0.1214(4)	3.3(4)
C42	0.0163(7)	0.6708(6)	0.0862(4)	5.0(5)
C43	−0.1067(8)	0.6372(7)	0.0561(5)	6.6(6)
C44	−0.1337(8)	0.5477(7)	0.0619(5)	7.0(6)
C45	−0.0493(9)	0.4863(7)	0.0963(5)	7.1(6)
C46	0.0748(8)	0.5175(6)	0.1261(4)	5.1(5)
C51	0.3744(7)	0.5580(5)	0.1530(4)	3.1(4)
C52	0.3928(7)	0.5353(5)	0.0915(4)	3.9(4)
C53	0.4628(8)	0.4583(6)	0.0823(4)	5.3(5)
C54	0.5183(9)	0.4064(6)	0.1359(5)	6.2(5)
C55	0.5056(9)	0.4280(6)	0.1981(4)	6.2(6)
C56	0.4333(8)	0.5045(6)	0.2078(4)	4.9(5)

Table 5. Crystal Data for Complexes 7a and 18

	7a	18
formula	C ₂₂ H ₂₁ F ₆ NOP ₂ Fe	C ₃₃ H ₃₁ Cl ₂ NP ₂ Pd
fw	547.2	680.75
cryst syst	triclinic	monoclinic
space group	<i>P</i> $\bar{1}$	<i>P</i> 2 ₁ / <i>n</i>
temp, K	298	298
<i>a</i> , Å	9.175(13)	10.460(3)
<i>b</i> , Å	17.514(9)	14.440(5)
<i>c</i> , Å	14.481(25)	20.558(8)
α , deg	90.51(8)	
β , deg	92.28(12)	101.99(3)
γ , deg	90.61(8)	
<i>V</i> , Å ³	2325(5)	3038(2)
2 θ range, deg	18.68–26.30	19.28–24.84
<i>F</i> (000)	1112	1384
<i>Z</i>	4	4
<i>D</i> _{calcd} , g cm ^{−3}	1.563	1.489
μ , cm ^{−1}	8.440	9.757
scan width	ω –2 θ	ω –2 θ
radiation	Mo K α	Mo K α
cryst dims, mm	0.20 × 0.30 × 0.35	0.30 × 0.30 × 0.35
scan width	2(1.00 + 0.35 tan θ)	2(0.90 + 0.35 tan θ)
transm range	0.551, 1.00	0.913, 1.00
2 θ _{max} , deg	45.0	45.0
no. of unique reflns	6055	3962
no. of reflns obsd ^a	1868	2821
computation	NRCSDP-VAX	NRCSDP-VAX
soln method	heavy atom	heavy atom
no. of params	475	352
<i>R</i>	0.089	0.044
<i>R</i> _w	0.084	0.033
<i>S</i>	2.67	2.51

^a $I > 2\sigma(I)$. ^b $R = \sum |F_o - F_c| / \sum (F_o)$; $R_w = [\sum w(F_o - F_c)^2 / \sum w(F_o)^2]^{1/2}$; $S = [\sum w(F_o - F_c)^2 / (\text{no. of reflns} - \text{no. of params})]$.

photochemical apparatus was a 450 W Conrad-Hanovia medium-pressure mercury lamp (Ace Glass).

All of the reaction, manipulation, and purification steps involving phosphines were performed under a dry nitrogen atmosphere. Tetrahydrofuran was distilled under nitrogen

from sodium benzophenone ketyl. Benzene and toluene were distilled from sodium under nitrogen. Dichloromethane was dried by CaH₂ and then distilled under nitrogen. Other chemicals and solvents were used from commercial sources without further purification.

N-(3-(Diphenylphosphino)propyl)triphenylphosphinimine, 1. To a benzene solution of 3-azidopropyl chloride (4.74 g, 39.6 mmol) was added a benzene solution of triphenylphosphine (10.4 g, 39.6 mmol). The resulting solution was stirred for 8 h, and the solvent was removed under reduced pressure. A white solid of (3-chloropropyl)tri-

Chart 1

phenylphosphinimine (15.14 g, 100%) was obtained, which was characterized by spectral methods: ^1H NMR δ 7.70–7.25 (m, 15 H, Ar H), 3.67 (t, J = 6.6 Hz, 2 H, $-\text{CH}_2\text{Cl}$), 3.20 (dt, J = 16, 6.4 Hz, 2 H, $-\text{CH}_2\text{NP}-$), 1.95 (dtt, J = 1.2, 6.4, 6.4 Hz, 2 H, $-\text{CH}_2-$); ^{13}C NMR δ (aliphatic carbon) 43.6 ($-\text{CCl}$), 42.1 (d, $J_{\text{P-C}}$ = 4.4 Hz, $-\text{CN}=\text{P}$), 37.8 (d, $J_{\text{P-C}}$ = 18.7 Hz, $-\text{CH}_2-$); ^{31}P NMR δ 12.4.

A solution of diphenylphosphide anion was prepared by the addition of a 1.6 M hexane solution of *n*-butyllithium (25 mL, 40 mmol) to a THF solution (100 mL) of diphenylphosphine (7.4 g, 39.7 mmol). This anion solution was then added to a solution of (3-chloropropyl)triphenylphosphinimine (15.4 g) in THF (100 mL) with stirring. The resulting mixture was kept stirred for 4 h, and solvents were removed under reduced pressure. The residue was dissolved in benzene and filtered. The filtrate was concentrated and recrystallized from acetonitrile to give the desired compound **1** as a white solid (13.2 g, 66%), which is quite sensitive to moisture: mp 120–122 °C; ^1H NMR (C_6D_6) δ 7.80–7.50 (m, 12 H, Ar H), 7.15–6.95 (m, 13 H, Ar H), 3.50 (dt, J = 16, 6.4 Hz, 2 H, $-\text{CH}_2\text{N}$), 2.41 (m, 2 H, $-\text{CH}_2\text{P}-$), 2.08 (m, 2 H, $-\text{CH}_2-$); ^{13}C NMR δ (aliphatic carbon) 46.8 (dd, J = 14.4, 4.6 Hz, $-\text{CNP}-$), 32.7 (dd, J = 20.9, 14.8 Hz, $-\text{CPPH}_2$), 26.4 (d, J = 11.5 Hz, $-\text{CH}_2-$); ^{31}P NMR (C_6D_6) δ 4.3, –16.0; ^{31}P NMR (CDCl_3) δ 12.1 (br), –15.0. HRMS Calcd for $\text{C}_{33}\text{H}_{31}\text{NP}_2$: m/z 503.1932. Found: m/z = 503.1933. Anal. Calcd for $\text{C}_{33}\text{H}_{31}\text{NP}_2$: C, 78.71; H, 6.21; N, 2.78. Found: C, 78.52; H, 6.26; N, 2.75.

[Ph₂P(CH₂)₃NPPH₃-P,N]Mo(CO)₄ (2). A mixture of **1** (201 mg, 0.40 mmol) and $\text{Et}_4\text{N}[\text{Mo}(\text{CO})_5\text{Br}]$ (175 mg, 0.38 mmol) in THF (10 mL) was heated to reflux for 8 h. The reaction mixture was filtered, and the filtrate was concentrated to give the desired complex **2** as a yellow solid (175 mg, 66%): mp 125–138 °C dec; IR (THF) ν_{CO} 2004, 1891, 1875, 1843 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.71–7.20 (m, 25 H, Ar H), 3.47 (dm, $J_{\text{P-H}}$ = 20.7 Hz, 2 H, $-\text{CH}_2\text{NP}-$), 2.62 (m, 2 H, $-\text{CH}_2\text{P}-$), 1.64 (m, 2 H, $-\text{CH}_2-$); ^{13}C NMR δ (aliphatic) 54.8 (d, J = 5.1 Hz, $-\text{CH}_2\text{NP}-$), 29.3 (d, J = 15.1 Hz, $-\text{CPPH}_2$), 28.0 (d, J = 8.6 Hz, $-\text{CH}_2-$); ^{31}P NMR δ 39.4 (d, J = 14 Hz), 16.3 (d, J = 14 Hz). Anal. Calcd for $\text{C}_{37}\text{H}_{31}\text{NO}_4\text{P}_2\text{Mo}$: C, 62.46; H, 4.39; N, 1.97. Found: C, 62.00; H, 4.80; N, 1.86.

[Ph₂P(CH₂)₃NPPH₃-P,N]W(CO)₄ (3). The preparation of **3** was similar to the procedure described for **2**. Complex **3** is a yellow solid: mp 179–186 °C dec; IR (THF) ν_{CO} 1998, 1881, 1863, 1838 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.22–7.40 (m, 25 H, Ar H), 3.67 (dm, $J_{\text{P-H}}$ = 19.5 Hz, 2 H, $-\text{CH}_2\text{NP}-$), 2.77 (m, 2 H, $-\text{CH}_2\text{P}-$), 1.67 (m, 2 H, $-\text{CH}_2-$); ^{13}C NMR δ (aliphatic) 56.1 (d, J = 5.7 Hz, $-\text{CNP}-$), 29.4 (d, J = 18.5 Hz, $-\text{CPPH}_2$), 27.8 (d, J = 8.3 Hz, $-\text{CH}_2-$); ^{31}P NMR δ 37.9 (d, J = 18 Hz), 2.67 (dt, $J_{\text{P-P}}$ = 18 Hz, $J_{\text{P-W}}$ = 224 Hz). Anal. Calcd for $\text{C}_{37}\text{H}_{31}\text{NO}_4\text{P}_2\text{W}$: C, 55.59; H, 3.91; N, 1.75. Found: C, 55.20; H, 3.90; N, 2.10.

General Procedure for Preparation of $\text{M}(\text{CO})_5(\text{CNCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2\text{-C})$ [$\text{M} = \text{Cr}, \text{Mo}, \text{W}$]. A mixture of $\text{M}(\text{CO})_6$ and **1** in equimolar quantity in THF was stirred at 25 °C for 40 h.

The reaction mixture was concentrated, and the residue was chromatographed on silica gel with elution of dichloromethane. A yellow band was collected and concentrated to give the desired product.

[Ph₂P(CH₂)₃NC-C]Cr(CO)₅ (4). A yellow liquid (46% yield): IR (CH_2Cl_2) ν_{CN} 2173 cm^{-1} , ν_{CO} 2066, 1955 cm^{-1} ; ^1H NMR δ 7.47–7.32 (m, 10 H, Ar H), 3.65 (t, J = 6.4 Hz, 2 H, $-\text{NCH}_2$), 2.18 (m, 2 H, $-\text{CH}_2\text{P}-$), 1.85 (m, 2 H, $-\text{CH}_2-$); ^{13}C NMR δ 216.9, 214.9, 162.5 (CN-), 137.5 (d, J = 11 Hz), 132.6 (d, J = 19 Hz), 129.0, 128.7 (d, J = 5.8 Hz), 45.3 (d, J = 14.3 Hz), 26.1 (d, J = 19.3 Hz, $-\text{CH}_2\text{P}-$), 25.0 (d, J = 12.7 Hz, $-\text{CH}_2-$); ^{31}P NMR δ –17.66. Anal. Calcd for $\text{C}_{21}\text{H}_{16}\text{NO}_5\text{PCr}$: C, 56.64; H, 3.62; N, 3.15. Found: C, 56.60; H, 3.10; N, 2.94.

[Ph₂P(CH₂)₃NC-C]Mo(CO)₅ (5). A yellow liquid (50%): IR (CH_2Cl_2) ν_{CN} 2173 cm^{-1} , ν_{CO} 2071, 1952 cm^{-1} ; ^1H NMR δ 7.51–7.33 (m, 10 H, Ar H), 3.67 (t, J = 6.5 Hz, 2 H, $-\text{NCH}_2$), 2.21 (m, 2 H, $-\text{CH}_2\text{PPh}_2$), 1.89 (m, 2 H, $-\text{CH}_2-$); ^{13}C NMR δ 206.7, 203.8, 153.4 (CN-), 137.4 (d, J = 12.5 Hz), 132.5 (d, J = 19 Hz), 129.0, 128.6 (d, J = 6.8 Hz), 44.8 (d, J = 14.3 Hz), 25.9 (d, J = 18.2 Hz), 24.9 (d, J = 13.7 Hz); ^{31}P NMR δ –17.71. Anal. Calcd for $\text{C}_{21}\text{H}_{16}\text{NO}_5\text{PMo}$: C, 51.55; H, 3.30; N, 2.80. Found: C, 51.52; H, 3.40; N, 2.83.

[Ph₂P(CH₂)₃NC-C]W(CO)₅ (6). A yellow liquid (67%): IR (CH_2Cl_2) ν_{CN} 2175 cm^{-1} , ν_{CO} 2068, 1956 cm^{-1} ; ^1H NMR δ 7.47–7.32 (m, 10 H, Ar H), 3.72 (t, J = 6.5 Hz, 2 H, $-\text{NCH}_2$), 2.19 (m, 2 H, $-\text{CH}_2\text{P}-$), 1.89 (m, 2 H, $-\text{CH}_2-$); ^{13}C NMR δ 196.1, 194.3, 143.0, 137.4 (d, J = 12.3 Hz), 132.6 (d, J = 18.2 Hz), 129.0, 128.7 (d, J = 6.9 Hz), 45.0 (d, J = 14.2 Hz), 26.10 (d, J = 18.3 Hz), 25.0 (d, J = 12.9 Hz); ^{31}P NMR δ –17.75. Anal. Calcd for $\text{C}_{21}\text{H}_{16}\text{NO}_5\text{PW}$: C, 43.70; H, 2.79; N, 2.43. Found: C, 43.77; H, 2.76; N, 2.44.

[Ph₂P(CH₂)₃NC-P,C]Fe(CO)₅ (7). A solution of $\text{CpFe}(\text{CO})_2\text{I}$ (75.9 mg, 0.25 mmol) and **1** (126 mg, 0.25 mmol) in toluene (20 mL) was stirred at room temperature for 4 h. A yellow precipitate formed during the reaction was collected by filtration. The yellow solid was washed with toluene and hexane to give the pure complex **7** (110 mg, 83%): mp 165–169 °C dec; IR (CHCl_3) 2089 (ν_{CN}), 1998 cm^{-1} (ν_{CO}); ^1H NMR (CDCl_3) δ 7.64–7.13 (m, 10 H), 4.97 (s, 5 H), 4.57–4.48 (m, 1 H), 3.67–3.56 (m, 1 H), 3.50–3.42 (m, 1 H), 2.74–2.62 (m, 1 H), 2.30–2.20 (m, 1 H), 2.20–2.05 (m, 1 H); ^{13}C NMR (CDCl_3) δ 213.0 (d, J = 22.3 Hz, $-\text{CO}$), 183.7 (d, J = 35.1 Hz, $-\text{NC}$), 133.9, 128.1 (m), 48.9, 32.0 (d, J = 31 Hz), 28.8 (d, J = 4.8 Hz); ^{31}P NMR (CDCl_3) δ 54.0; FAB m/z = 402. Anal. Calcd for $\text{C}_{22}\text{H}_{21}\text{INOFe}$: C, 49.94; H, 4.00; N, 2.65. Found: C, 49.59; H, 4.05; N, 2.53.

[Ph₂P(CH₂)₃NC-P,C]Fe(CO)₅PF₆ (7a). A solution of **7** and excess NH_4PF_6 in acetone was stirred for 4 h. The reaction mixture was filtered and concentrated, and the residue was recrystallized from a mixed solvent (hexane/ CH_2Cl_2) to give **7a** as a yellow solid quantitatively: mp 175–180 °C dec; IR (KBr) 2098 (ν_{CN}), 2011 cm^{-1} (ν_{CO}); ^1H NMR (CDCl_3) δ 7.64–7.13 (m, 10 H, Ar H), 4.87 (s, 5 H, Cp H), 4.03 (m, 1 H), 3.50

(m, 1 H), 2.96 (m, 1 H), 2.60 (m, 1 H), 2.20 (m, 1 H), 2.00 (m, 1 H); ^{31}P NMR (CDCl_3) δ 54.0, -143.6 (PF_6).

[$\{\text{Ph}_2\text{P}(\text{CH}_2)_3\text{NH}(n\text{-PrNH})\text{C}=\text{Fe}(\text{CO})\}\text{I}$ (8). A mixture of *n*-propylamine (5 mL) and **7** (450 mg, 0.166 mmol) in dichloromethane (20 mL) was stirred at 25 °C for 1 h. On removal of the solvent, the residue was crystallized from chloroform and pentane to give **8** as a yellow crystalline solid (435 mg, 87%): mp 185–190 °C dec; IR (CH_2Cl_2) $\nu(\text{CO})$ 1949 cm^{-1} ; ^1H NMR (CDCl_3) δ 8.31 (br, 1 H), 7.49–7.12 (m, 10 H), 6.40 (br, 1 H), 4.53 (s, 5 H), 4.35 (m, 1 H), 3.60 (m, 2 H), 3.50 (m, 1 H), 2.79 (m, 2 H), 2.17 (m, 1 H), 2.11 (m, 2 H), 1.30 (m, 1 H), 1.11 (t, $J = 7$ Hz, 3 H); ^{13}C NMR (CDCl_3) δ 218.6 (d, $J = 31.8$ Hz), 209.8 (d, $J = 25.6$ Hz), 139.2 (d, $J = 48.3$ Hz), 132.9 (d, $J = 47$ Hz), 132.8 (d, $J = 9.4$ Hz), 131.02 (d, $J = 1.9$ Hz), 130.2 (d, $J = 1.7$ Hz), 130.0 (d, $J = 8.6$ Hz), 129.0 (d, $J = 9.5$ Hz), 128.6 (d, $J = 9.8$ Hz), 84.7, 48.2, 45.5, 35.5 (d, $J = 22$ Hz), 24.8, 21.9, 11.6; ^{31}P NMR (CDCl_3) δ 56.5. Anal. Calcd for $\text{C}_{25}\text{H}_{30}\text{IN}_2\text{POFe}$: C, 51.05; H, 5.14; N, 4.76. Found: C, 52.12; H, 5.07; N, 4.84.

[$\text{CpRu}\{\text{Ph}_2\text{P}(\text{CH}_2)_3\text{NPPH}_3\text{-P}\}(\text{PPh}_3)\text{Cl}$ (9). A mixture of $\text{CpRu}(\text{PPh}_3)_2\text{Cl}$ (634 mg, 0.873 mmol) and **1** (440 mg, 0.873 mmol) in benzene (40 mL) was heated to reflux for 4 h. The benzene solvent was replaced by ether, and a yellow precipitate was formed and collected. The precipitate was subsequently washed with ether (20 mL $\times$ 2) to give complex **5** as a yellow solid (570 mg, 68%): mp 122–125 °C dec; ^1H NMR (CDCl_3) δ 7.80–6.95 (m, 40 H), 4.03 (s, 5 H), 2.61 (dt, $J = 20$, 7.8 Hz, 2 H), 2.21–2.11 (m, 1 H), 1.38–1.20 (m, 1 H), 0.80–0.60 (m, 2 H); ^{13}C NMR δ (aliphatic) 80.5, 43.0 (d, $J = 11.6$ Hz), 28.4 (d, $J = 8.9$ Hz), 18.9 (dd, $J = 38.7$, 20 Hz); ^{31}P NMR δ 44.1 (d, $J = 41.4$ Hz), 36.8 (d, $J = 41.4$ Hz), 13.0 (br); FAB ($m + 1$)/ $z = 968$. Anal. Calcd for $\text{C}_{56}\text{H}_{51}\text{NP}_3\text{RuCl}$: C, 69.52; H, 5.31; N, 1.45. Found: C, 69.31; H, 5.24; N, 1.57.

[$\text{CpRu}\{\text{Ph}_2\text{P}(\text{CH}_2)_3\text{NC-C}\}(\text{CO})\text{I}$ (10). To a solution of $\text{CpRu}(\text{CO})_2\text{I}$ (200 mg, 0.57 mmol) in dichloromethane was added compound **1** (302 mg, 0.60 mmol); the resulting mixture was stirred at room temperature for 10 h. Removal of solvent yielded a liquid residue, which was chromatographed on alumina (10 g) with elution with dichloromethane. A red band eluate was collected and concentrated to give the complex **6** as a red liquid (272 mg, 83%): IR (CH_2Cl_2) $\nu(\text{NC})$ 2170, $\nu(\text{CO})$ 1973 cm^{-1} ; ^1H NMR δ 7.46–7.24 (m, 10 H), 5.14 (s, 5 H), 3.82 (t, $J = 6.4$ Hz, 2 H), 2.25–2.19 (m, 2 H), 1.92–1.79 (m, 2 H); ^{13}C NMR δ 199.5 (CO), 145.1 (CN), 137.5–128.6 (Ar C), 84.4 (Cp-), 45.7 (d, $J = 14.5$ Hz), 26.3 (d, $J = 18.1$ Hz), 24.8 (d, $J = 11.9$ Hz); ^{31}P NMR δ -17.3. Anal. Calcd for $\text{C}_{22}\text{H}_{21}\text{INOPRu}$: C, 46.01; H, 3.69; N, 2.44. Found: C, 45.78; H, 3.69; N, 2.46.

[$\text{Cp}(\text{PPh}_3)\text{ClRu}\{\mu\text{-Ph}_2\text{P}(\text{CH}_2)_3\text{NC-P-C}\}\text{RuCp}(\text{CO})\text{I}$ (11). **Method 1.** A mixture of **9** (140 mg, 0.144 mmol) and $\text{CpRu}(\text{CO})_2\text{I}$ (50.4 mg, 0.144 mmol) in toluene (10 mL) was stirred at room temperature for 10 h. Concentration of the reaction mixture gave the crude product as a red solid, which was purified by chromatography on alumina with elution of dichloromethane. Recrystallization from CHCl_3 /pentane at -20 °C provided the desired complex **11** as an orange crystalline solid (107 mg, 72%).

Method 2. A benzene solution of **10** (60 mg, 0.104 mmol) and $\text{CpRu}(\text{PPh}_3)_2\text{Cl}$ (75.8 mg, 0.104 mmol) was heated to reflux for 4 h. Replacement of benzene with ether solvent yielded a red precipitate solid, which was washed with ether (20 mL, $\times$ 2) to provide pure **11** as an orange-red solid (63 mg, 58%).

Both methods provided formation of diastereomeric pairs of **11**, which could not be separated at present. IR (CH_2Cl_2): $\nu(\text{NC})$ 2170, $\nu(\text{CO})$ 1970 cm^{-1} . ^1H NMR: δ 7.90–7.88 (m, 2 H), 7.50–7.04 (m, 23 H), 5.14 (s, 5 H), 4.13 (s, 5 H), 3.31–3.13 (m, 2 H), 2.31–2.22 (m, 1 H), 1.29–1.21 (m, 1 H), 1.10–1.00 (m, 1 H), 0.88–0.74 (m, 1 H). ^{31}P NMR: δ 44.12 (d, $J = 41.4$ Hz), 35.74 (d, $J = 41.4$ Hz) as one pair of diastereomers; δ 44.05 (d, $J = 41.4$ Hz), 35.78 (d, $J = 41.4$ Hz). Anal. Calcd for $\text{C}_{45}\text{H}_{41}\text{ClINOP}_2\text{Ru}_2\text{O} \cdot 0.5\text{CHCl}_3$: C, 49.77; H, 3.81; N, 1.24. Found: C, 49.28; H, 3.74; N, 1.24.

$\text{Br}(\text{CO})_4\text{Re}(\text{CNCH}_2\text{CH}_2\text{PPh}_2\text{-C})$ (12). A mixture of $\text{Re}(\text{CO})_5\text{Br}$ (73 mg, 0.18 mmol) and **1** (90.5 mg, 0.18 mmol) in CH_2Cl_2 (10 mL) was heated to reflux for 2 h. After removal of solvent, the residue was chromatographed on silica gel with dichloromethane as the eluent. The eluate was collected and concentrated to give the desired complex **12** as a colorless liquid (75 mg, 66%): IR (CH_2Cl_2) ν_{CN} 2222 cm^{-1} , ν_{CO} 2112, 2020, 1958 cm^{-1} ; ^1H NMR δ 7.47–7.30 (m, 10 H, Ar H), 3.87 (t, $J = 6.5$ Hz, 2 H, $-\text{NCH}_2-$), 2.22 (m, 2 H, $-\text{PCH}_2-$), 1.93 (m, 2 H, $-\text{CH}_2-$); ^{13}C NMR δ 182.3, 181.1, 180.1 ($-\text{CO}$), 137.2 (d, $J = 11.2$ Hz), 132.6 (d, $J = 19.2$ Hz), 129.0, 128.6 (d, $J = 6.8$ Hz) (Ar C), 129.2 ($-\text{CN}$), 45.4 (d, $J = 15.1$ Hz, $-\text{NCH}_2-$), 25.6 (d, $J = 18.7$ Hz, $-\text{PCH}_2-$), 24.7 (d, $J = 13.2$, $-\text{CH}_2-$); ^{31}P NMR δ -17.83. Anal. Calcd for $\text{C}_{20}\text{H}_{16}\text{BrNPO}_4\text{Re}$: C, 38.04; H, 2.22; N, 2.55. Found: C, 37.95; H, 2.31; N, 2.52.

$\text{Br}(\text{CO})_3\text{Re}(\text{CNCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2\text{-P,C})$ (13). A solution of **12** (50 mg, 0.08 mmol) in benzene (10 mL) was heated to reflux for 4 h. The reaction mixture was chromatographed on silica gel (5 g) with elution with CH_2Cl_2 . The eluate was collected and concentrated to give **13** as a clear colorless liquid (35 mg, 73%): IR ν_{CN} 2137 cm^{-1} , ν_{CO} 2039, 1975, 1917 cm^{-1} ; ^1H NMR δ 7.68–7.36 (m, 10 H), 4.00 (m, 1 H), 3.44 (m, 2 H), 2.76 (m, 1 H), 2.07 (m, 2 H); ^{13}C NMR δ 190.2 (d, $J = 8.9$ Hz), 187.83 (d, $J = 58.5$ Hz), 187.78 (d, $J = 6.2$ Hz) ($-\text{CO}$), 163.0 (d, $J = 6.8$ Hz, $-\text{CN}$), 134.3–128.0 (Ar C), 47.2, 28.7 (d, $J = 2.7$ Hz), 27.7 (d, $J = 28$ Hz); ^{31}P NMR δ -4.33. Anal. Calcd for $\text{C}_{19}\text{H}_{16}\text{NPO}_3\text{ReBr}$: C, 37.82; H, 2.67; N, 2.32. Found: C, 37.57; H, 2.73; N, 2.30.

$(\text{CO})_3\text{BrRe}(\text{CNCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2\text{-C})_2$ (14). A mixture of **1** (920 mg, 1.827 mmol) and $\text{BrRe}(\text{CO})_5$ (370 mg, 0.911 mmol) in benzene (40 mL) was stirred at 25 °C for 10 h. After concentration of the reaction mixture, the residue was chromatographed on silica gel with dichloromethane as the eluent. The eluate was concentrated to give **14** as a colorless liquid (565 mg, 73%): IR (CHCl_3) ν_{CN} , 2041, 1981, 1926 cm^{-1} (ν_{CO}); ^1H NMR δ 7.47–7.31 (m, 20 H, Ar H), 3.68 (t, $J = 6.4$ Hz, 4 H, CNCH_2-), 2.17 (m, 4 H, $-\text{CH}_2\text{PPh}_2$), 1.83 (m, 4 H, $-\text{CH}_2-$); ^{13}C NMR δ 186.1, 183.8, 137.1 (d, $J = 8.6$ Hz), 137.0 (d, $J = 8.9$ Hz), 134.3 ($-\text{CN}$), 132.4 (d, $J = 6.8$ Hz), 132.2 (d, $J = 6.6$ Hz), 128.7, 128.6, 128.3 (d, $J = 6.6$ Hz), 44.6 (d, $J = 14.5$ Hz), 25.3 (d, $J = 18.5$ Hz), 24.2 (d, $J = 13.1$ Hz); ^{31}P NMR δ -17.80; FABMS ($m + 1$)/ $z = 857.0$. Anal. Calcd for $\text{C}_{35}\text{H}_{32}\text{N}_2\text{P}_2\text{O}_3\text{ReBr}$: C, 49.07; H, 3.76; N, 3.27. Found: C, 49.01; H, 3.80; N, 3.26.

$(\text{CO})_5\text{Re}_2(\text{CNCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2\text{-C})$ (15). A solution of **1** (42 mg, 0.083 mmol) and $\text{Re}_2(\text{CO})_{10}$ (54 mg, 0.083 mmol) in toluene (10 mL) was stirred at 25 °C for 10 h. The reaction mixture was chromatographed on silica gel with dichloromethane as the eluent. Upon concentration of the eluate, the desired product was obtained as a colorless liquid (67 mg, 92%): IR 2187 (ν_{CN}), 2099, 2074, 2046, 2008, 1994, 1968, 1942 cm^{-1} (ν_{CO}); ^1H NMR δ 7.50–7.33 (m, 10 H), 3.85 (t, $J = 6.7$ Hz, 2 H), 2.12 (m, 2 H), 1.85 (m, 2 H); ^{13}C NMR δ 194.6, 192.0, 184.6 ($-\text{CO}$), 134.5 ($-\text{CN}$), 137.3 (d, $J = 12$ Hz), 132.6 (d, $J = 19.3$ Hz), 129.0 128.7 (d, $J = 7.6$ Hz) (Ar C), 45.6 (d, $J = 14.4$ Hz), 25.8 (d, $J = 19.5$ Hz), 25.0 (d, $J = 13.9$ Hz); ^{31}P NMR δ -17.5. Anal. Calcd for $\text{C}_{25}\text{H}_{16}\text{NO}_9\text{Re}_2$: C, 34.21; H, 1.84; N, 1.60. Found: C, 34.32; H, 1.80; N, 1.59.

$(\text{CO})_4\text{Re}\{\mu_2\text{-CNCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2\text{-P,C}\}\text{Re}(\text{CO})_4$ (16). A solution of **15** (640 mg, 0.729 mmol) in chloroform was irradiated with light for 2 h. After concentration, the residue was chromatographed on silica gel with dichloromethane as eluent. The eluate was concentrated to give the desired product **16** as a white solid (450 mg, 73%): mp 165–185 °C dec; IR (KBr) ν 2156 ($-\text{CN}$), 2081, 2062, 2034, 1973, 1958 cm^{-1} ($-\text{CO}$); ^1H NMR δ 7.64–7.47 (m, 10 H), 3.86 (t, $J = 6$ Hz, 2 H), 2.92 (m, 2 H), 1.83 (m, 2 H); ^{13}C NMR δ 185.4 (d, $J = 9$ Hz), 183.5, 183.3 (d, $J = 7$ Hz), 182.8, 182.1, 180.6, 178.9 (CN), 132.3, 132.2 (d, $J = 10.5$ Hz), 131.5 (d, $J = 2$ Hz), 129.4 (d, $J = 9.7$ Hz), 45.2 (d, $J = 17$ Hz), 23.5, 22.9 (d, $J = 30$ Hz); ^{31}P NMR δ -0.3. Anal. Calcd for $\text{C}_{24}\text{H}_{16}\text{NO}_8\text{Re}_2$: C, 35.21; H, 1.97; N, 1.71. Found: C, 34.93; H, 2.05; N, 1.69.

Br(CO)₄Re{μ₂-CNCH₂CH₂CH₂PPh₂-P,C}Re(CO)₄Br (17). Bromine (31 mg, 0.194 mmol) was added to a solution of **16** (140 mg, 0.166 mmol) in dichloromethane (100 mL) at room temperature. After stirring for 3 h, the reaction mixture was concentrated and the residue was chromatographed on silica gel with ethyl acetate and hexane (1/4) as the eluent. The eluate was collected and concentrated to give **17** as a clear, colorless liquid (125 mg, 75%): IR (CH₂Cl₂) 2221 cm⁻¹ (–CN), 2110, 2021, 1954 cm⁻¹ (–CO); ¹H NMR δ 7.62–7.47 (m, 10 H), 3.88 (t, *J* = 6 Hz, 2 H), 3.02 (m, 2 H), 1.80 (m, 2 H); ¹³C NMR δ 184.2 (d, *J* = 9.2 Hz), 182.5 (d, *J* = 6.6 Hz), 182.3, 181.7, 181.0, 179.9, 132.0 (d, *J* = 10.6 Hz), 131.4, 131.3, 130.7, 129.1 (d, *J* = 9.8 Hz), 44.8 (d, *J* = 17 Hz), 24.1 (d, *J* = 31.3 Hz), 23.2; ³¹P NMR δ –5.12. Anal. Calcd for C₂₄H₁₆Br₂NO₈Re₂: C, 29.46; H, 1.65; N, 1.43. Found: C, 29.31; H, 1.75; N, 1.42.

Cl₂Pd(Ph₂PCH₂CH₂CH₂NPPh₃-P,N) (18). A mixture of **1** (106 mg, 0.21 mmol) and Pd(COD)Cl₂ (60 mg, 0.21 mmol) in dichloromethane (15 mL) was stirred at 25 °C for 8 h. On removal of solvents, the residue was dissolved in chloroform and pentane and the desired complex **18** crystallized as a yellow-orange solid (136 mg, 95%): mp 154–158 °C dec; ¹H NMR δ 7.67–7.33 (m, 25 H), 3.21 (m, 2 H), 2.05 (m, 2 H), 1.86 (m, 2 H); ¹³C NMR δ (aliphatic) 45.5 (d, *J* = 2.9 Hz), 25.3 (d, *J* = 9.5 Hz), 22.5 (d, *J* = 33.5 Hz); ³¹P NMR δ 40.5 (d, *J* = 3.3

Hz), 17.2 (d, *J* = 3.3 Hz). Anal. Calcd for C₃₃H₃₁Cl₂NP₂Pd: C, 58.21; H, 4.59; N, 2.06. Found: C, 58.19; H, 4.84; N, 1.81.

X-ray Crystallography. Single crystals suitable for X-ray analysis of complexes **7a** and **18** were obtained in each case by slow evaporation of a chloroform/pentane solution under air. Cell parameters were determined on a CAD-4 diffractometer at 298 K by a least-squares treatment. Atomic scattering factors were taken from ref 24. The NRCC SDP VAX program was used for calculation.²⁵ Crystal data of these complexes are summarized in Table 5, and their non-hydrogen atomic coordinates are listed in Tables 2 and 4, respectively.

Acknowledgment. We thank the National Science Council of the Republic of China for grant support (NSC84-2113-M002-018).

Supplementary Material Available: Anisotropic thermal parameters and complete bond distances and bond angles for **7a** and **18** (7 pages). Ordering information is given on any current masthead page.

OM940875K

(24) *International Tables for X-ray Crystallography*; Kynoch Press: Birmingham, U.K., 1974; Vol. IV.

(25) Gabe, E. J.; Lee, F. L. *Acta Crystallogr.* **1981**, A37, S339.