Two [2]pseudorotaxane-like complexes and their corresponding [2]rotaxanes stabilized via interactions on opposite ends of the same macrocycle

Wei-Chung Hung^a, Liang-Yun Wang^a, Chien-Chen Lai^b, Yi-Hung Liu^a,
Shie-Ming Peng^a, Sheng-Hsien Chiu^{a,*}

^aDepartment of Chemistry, National Taiwan University, No. 1, Sec. 4, Roosevelt Road, Taipei 10617, Taiwan, ROC

^bInstitute of Molecular Biology, National Chung Hsing University and Department of Medical Genetics, China Medical University Hospital, Taichung, Taiwan, ROC

ARTICLE INFO

Article history:

Received 21 August 2008

Revised 10 October 2008

Accepted 10 October 2008

Available online 7 November 2008

ABSTRACT

A multiple-use macrocycle recognizes dibenzylammonium ions and 2,6-lutidine derivatives, each in a [2]pseudorotaxane-like manner, through interactions with its diethylene glycol (hydrogen bonding) and 2,6-pyridinedicarboxamide (Pd²⁺ chelation) spacers, respectively. We characterized these complexes in the solid state (X-ray crystallography) and in solution (¹H NMR spectroscopy). The synthesis of two corresponding [2]rotaxanes confirmed that these recognition systems possess [2]pseudorotaxane geometries in solution.

© 2008 Elsevier Ltd. All rights reserved.

Rotaxanes and catenanes attract a great deal of attention because of their potential for applications in mesoscale devices and molecular machinery on the nanoscale.¹ The synthesis of these interlocked systems usually requires the recognition of a macrocyclic unit by a recognition unit on a threadlike component.² For the realization of interlocked molecular switches and actuators, however, more than one type of recognition site is generally needed to allow itinerant migration of the macrocycle(s) between different recognition sites under the influence of an external stimulus; this binding must not only be discriminative but also sufficiently strong at each of these recognition stations.³ Nevertheless, macrocycles that can recognize more than one type of guest in a pseudorotaxane-like manner in solution with reasonable binding affinity are relatively rare. The crown ethers dibenzo[24]crown-8 (DB24C8) and bis-*p*-phenylene[34]crown-10 (BPP34C10) are particularly notable examples that have been used to construct several molecular switches because they can recognize several guests—dibenzylammonium ions (DBA⁺),⁴ 1,2-bis(pyridinium)ethane units,⁵ or bipyridinium dications⁶—in [2]pseudorotaxane-like geometries in solution with sufficiently high binding affinities. Intuitively, dual- or multiple-use macrocycles for future supramolecular catalytic systems or for the preparation of sensitive molecular switches can be developed by linking two or more different recognition systems as loops within a single macrocycle. The design of such a system requires a delicate balance in the molecular structure to ensure that both recognition systems operate independently. Previously, we reported that macrocycle **1**, which possesses two xylol

rings linked by diethylene glycol and 2,6-pyridinedicarboxamide spacers, acts as a host molecule that complexes diphenylurea derivatives through the cooperative interactions with its two opposing recognition units.⁷ Herein, we report that the opposing diethylene glycol and 2,6-pyridinedicarboxamide units of this macrocycle also act independently to recognize both the DBA⁺ ion and 2,6-lutidine in solution.

In an earlier study, we found that the recognition of a DBA⁺ ion requires a macrocycle featuring only one diethylene glycol chain and two phenolic rings.⁸ Because the 2,6-pyridinedicarboxamide motif places the two phenolic rings a suitable distance apart for

Scheme 1.

* Corresponding author.

E-mail address: shchiu@ntu.edu.tw (S.-H. Chiu).

the binding of a guest,⁹ we suspected that macrocycle **1** (Scheme 1) would form a pseudorotaxane-like complex with the DBA⁺ ion, stabilized primarily through [N⁺–H...O] and [N⁺C–H... π] interactions.

The ¹H NMR spectrum of an equimolar (10 mM) mixture of macrocycle **1** and DBA·PF₆ in CD₃NO₂ at room temperature displays changes in the chemical shifts of the protons of the complex relative to those of its free components (Fig. 1); these time-averaged signals suggest that the rates of complexation and decomplexation of **1** and DBA⁺ are rapid under these conditions. The spectrum of the complex reveals an upfield shift of the signal for the protons of the benzylic methylene groups adjacent to the NH₂⁺ center of the DBA⁺ ion as well as a separation of the ‘tight’ multiplet (δ 3.52–3.62) for the methylene protons of the OCH₂–CH₂O units of the free macrocycle into two signals (δ 3.41 and δ 3.62) for the complex; these features suggest the existence of [N⁺–H...O] and [N⁺C–H... π] hydrogen bonds between macrocycle **1** and the DBA⁺ ion.⁸ Accordingly, we suspected the [2]pseudorotaxane-like assembly [**1**⊃DBA⁺] (Scheme 1) to be the likely structure of the complex formed between **1** and the DBA⁺ ion in CD₃NO₂. From a ¹H NMR spectroscopic dilution experiment, we determined the association constant (*K*_a) for this complex in CD₃NO₂ to be 330 ± 30 M^{−1}.¹⁰

We grew single crystals suitable for X-ray crystallography through liquid diffusion of *i*Pr₂O into a CH₃NO₂ solution of macrocycle **1** and DBA·PF₆. The solid state structure^{11,12} (Fig. 2) reveals a [2]pseudorotaxane-like molecular geometry for the complex [**1**⊃DBA⁺]; it also indicates that the complex is stabilized through both [N⁺–H...O] hydrogen bonds and [N⁺C–H... π] interactions (the distances between the phenolic rings’ centroids and the sandwiched methylene carbons are both around 3.6 Å).

This result confirms our previous finding⁸ that, with suitable design, weak noncovalent interactions can be harnessed to play extremely important roles in stabilizing macrocycle/dialkylammonium ion complexes.

To confirm the existence of the [2]pseudorotaxane [**1**⊃DBA⁺] in solution, we constructed a [2]rotaxane based on this recognition system. Adding triethyl phosphite (200 mM) to a solution of the benzylic azide **2**·H·PF₆ (100 mM) and macrocycle **1** (150 mM) in CH₂Cl₂ gave the corresponding rotaxane, which dissociated slowly during the purification process, presumably because the stoppering groups were not bulky enough.¹³ Indeed, when we added tributyl phosphite under otherwise identical conditions, we isolated

Figure 1. Partial ¹H NMR spectra (400 MHz, CD₃NO₂, 298 K) of (a) macrocycle **1**, (b) an equimolar mixture of **1** and DBA·PF₆ (10 mM), and (c) DBA·PF₆.

Figure 2. (a) Top and (b) side views of the solid state structure of the [2]pseudorotaxane [**1**⊃DBA⁺]. The hydrogen bonding geometries, D...A, H...A [Å], and D–H...A [°]: (a) 2.98, 2.11, 168.9; (b) 2.88, 2.00, 174.1; (c) 3.64, 2.97, 135.5; (d) 3.72, 2.75, 152.0. D = hydrogen bond donor, A = hydrogen bond acceptor.

the [2]rotaxane **3**·H·PF₆ in 23% yield (Scheme 2).¹⁴ After dissolving the phosphoramidate-stoppered [2]rotaxane **3**·H·PF₆ in CD₃SOCD₃, we observed no signals for the free macrocycle **1** in the ¹H NMR spectrum, confirming the interlocked nature of its components (Fig. 3). To test the thermal stability of **3**·H·PF₆, we heated this

Scheme 2.

Figure 3. Partial ^1H NMR spectra (400 MHz, CD_3SOCD_3 , 298 K) of (a) the [2]rotaxane **3**-H- PF_6 and (b) macrocycle **1**. #: The signal of H_2O .

solution at 323 K while monitoring its ^1H NMR spectra. Again, we did not observe any signals for the free macrocycle within 2 h, suggesting that the dibutyl phosphoramidate groups are stoppers that can mechanically interlock the macrocycle **1** along the rodlike component of the dumbbell-shaped moiety.

Having proved that the diethylene glycol unit appended to the two aromatic rings in **1** was capable of complexing DBA^+ ions in CD_3NO_2 , we turned our attention to the complexing behavior of the opposing 2,6-pyridinedicarboxamide moiety, which is known to complex lutidine units via chelation of palladium(II).¹⁵ Thus, we treated macrocycle **1** with $\text{Pd}(\text{OAc})_2$ in CH_3CN for 3 h at room temperature to afford (Scheme 3) the corresponding Pd(II) complex **4** (75% yield), which we then mixed with 2,6-lutidine in CHCl_3 to produce the Pd(II) complex **5** in quantitative yield. We obtained single crystals suitable for X-ray crystallography upon liquid diffu-

sion of $i\text{Pr}_2\text{O}$ into a CH_3CN solution of complex **5**. The solid state structure¹⁶ reveals (Fig. 4) a [2]pseudorotaxane-like molecular geometry in which the Pd(II) atom holds the 2,6-lutidine and 2,6-pyridinedicarboxamide units together with their aromatic planes aligned perpendicularly.

To construct a [2]rotaxane based on this recognition system, we reacted the diol **6** with the Pd(II) complex **4** in CHCl_3 to generate the [2]pseudorotaxane complex **7** presenting hydroxyl groups at its termini. Reacting complex **7** with the bulky isocyanate **8** in the presence of di-*n*-butyltin dilaurate in CH_2Cl_2 at room temperature led us to isolate the [2]rotaxane **9** in 83% yield after column chromatography (Scheme 3).¹⁷ The successful preparation of this [2]rotaxane suggests that the 2,6-pyridinedicarboxamide unit within macrocycle **1** is capable of recognizing pyridine derivatives such as **6** in a pseudorotaxane-like manner in solution under the templating influence of Pd(II).¹⁸

Scheme 3.

Figure 4. (a) Top and (b) side views of the solid state structure of complex **5**.

Macrocyclic **1** is a multiple-use macrocycle that recognizes diphenylurea, DBA⁺, and 2,6-lutidine derivatives through interactions with either its diethylene glycol or Pd²⁺-chelated 2,6-pyridine diamide moieties. The solid state structure of [1⊃DBA⁺] confirmed that [N⁺C–H⋯π] interactions play an important role in stabilizing this complex. We aim to use multiple-guest-binding hosts such as **1** as components of future molecular catalytic systems and controllable molecular switches.

Acknowledgment

We thank the National Science Council for financial support (NSC 95-2113-M-002-016-MY3).

References and notes

- (a) Collier, C. P.; Mattersteig, G.; Wong, E. W.; Luo, Y.; Beverly, K.; Sampaio, J.; Raymo, F. M.; Stoddart, J. F.; Heath, J. R. *Science* **2000**, *289*, 1172–1175; (b) Saha, S.; Leung, K. C.-F.; Nguyen, T. D.; Stoddart, J. F.; Zink, J. I. *Adv. Funct. Mater.* **2007**, *17*, 685–693; (c) Green, J. E.; Choi, J. W.; Boukai, A.; Bunimovich, Y.; Johnston-Halperin, E.; Delonno, E.; Luo, Y.; Sheriff, B. A.; Xu, K.; Shin, Y. S.; Tseng, H.-R.; Stoddart, J. F.; Heath, J. R. *Nature* **2007**, *445*, 414–417.
- Molecular Catenanes Rotaxanes and Knots*; Sauvage, J.-P., Dietrich-Buchecker, C., Eds.; VCH-Wiley: Weinheim, 1999.
- (a) *Molecular Switches*; Feringa, B. L., Ed.; Wiley-VCH: Weinheim, 2001; (b) Elizarov, A. M.; Chiu, S.-H.; Stoddart, J. F. *J. Org. Chem.* **2002**, *67*, 9175–9181; (c) Huang, F.; Switek, K. A.; Gibson, H. W. *Chem. Commun.* **2005**, 3655–3657.
- (a) Fyfe, M. C. T.; Stoddart, J. F. *Adv. Supramol. Chem.* **1999**, *5*, 1–53; (b) Cantrill, S. J.; Pease, A. R.; Stoddart, J. F. *J. Chem. Soc., Dalton Trans.* **2000**, 3715–3734; (c) Clifford, T.; Abushamleh, A.; Busch, D. H. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99*, 4830–4836; (d) Gibson, H. W.; Yamaguchi, N.; Jones, J. W. *J. Am. Chem. Soc.* **2003**, *125*, 3522–3533; (e) Arico, F.; Badjic, J. D.; Cantrill, S. J.; Flood, A. H.; Leung, K. C.-F.; Liu, Y.; Stoddart, J. F. *Top. Curr. Chem.* **2005**, *249*, 203–259; (f) Griffiths, K. E.; Stoddart, J. F. *Pure Appl. Chem.* **2008**, *80*, 485–506.
- (a) Loeb, S. J.; Wisner, J. A. *Chem. Commun.* **1998**, 2757–2758; (b) Hoffart, D. J.; Loeb, S. J. *Angew. Chem., Int. Ed.* **2005**, *44*, 901–904; (c) Vella, S. J.; Tiburcio, J.; Loeb, S. J. *Chem. Commun.* **2007**, 4752–4754.
- (a) Allwood, B. L.; Spencer, N.; Shahriari-Zavareh, H.; Stoddart, J. F.; Williams, D. J. *J. Chem. Soc., Chem. Commun.* **1987**, 1064–1066; (b) Asakawa, M.; Ashton, P. R.; Ballardini, R.; Balzani, V.; Belohradsky, M.; Gandolfi, M. T.; Kocian, O.; Prodi, L.; Raymo, F. M.; Stoddart, J. F.; Venturi, M. *J. Am. Chem. Soc.* **1997**, *119*, 302–310; (c) Miljanic, O. S.; Stoddart, J. F. *Proc. Natl. Acad. Sci. U.S.A.* **2007**, *104*, 12966–12970; (d) Huang, F.; Jones, J. W.; Gibson, H. W. *J. Org. Chem.* **2007**, *72*, 6573–6576; (e) Wang, F.; Han, C.; He, C.; Zhou, Q.; Zhang, J.; Wang, C.; Li, N.; Huang, F. *J. Am. Chem. Soc.* **2008**, *130*, 11254–11255.
- Huang, Y.-L.; Hung, W.-C.; Lai, C.-C.; Liu, Y.-H.; Peng, S.-M.; Chiu, S.-H. *Angew. Chem., Int. Ed.* **2007**, *46*, 6629–6633.
- (a) Cheng, P.-N.; Hung, W.-C.; Chiu, S.-H. *Tetrahedron Lett.* **2005**, *46*, 4239–4242; (b) Cheng, P.-N.; Huang, P.-Y.; Li, W.-S.; Ueng, S.-H.; Hung, W.-C.; Liu, Y.-H.; Lai, C.-C.; Peng, S.-M.; Chao, I.; Chiu, S.-H. *J. Org. Chem.* **2006**, *71*, 2373–2383; (c) Cheng, K.-W.; Lai, C.-C.; Chiang, P.-T.; Chiu, S.-H. *Chem. Commun.* **2006**, 2854–2856; (d) Chiu, C.-W.; Lai, C.-C.; Chiu, S.-H. *J. Am. Chem. Soc.* **2007**, *129*, 3500–3501; (e) Hsueh, S.-Y.; Cheng, K.-W.; Lai, C.-C.; Chiu, S.-H. *Angew. Chem., Int. Ed.* **2008**, *47*, 4436–4439.
- Bisson, A. P.; Lynch, V. M.; Monahan, M.-K. C.; Anslyn, E. V. *Angew. Chem., Int. Ed.* **1997**, *36*, 2340–2342.
- Connors, K. A. *Binding Constants*; Wiley: New York, 1987.
- CCDC 703716 and 703717 contain the supplementary crystallographic data for [1⊃DBA][PF₆] and **5**, respectively. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- Crystal data for [1⊃DBA][DBA][2PF₆]: [C₅₅H₆₁N₅O₅][PF₆]₂; *M*_r = 1162.03; triclinic; space group *P*1; *a* = 11.1624(4); *b* = 13.2258(5); *c* = 19.4664(7) Å; *V* = 2797.18(18) Å³; *ρ*_{calcd} = 1.380 g cm^{−3}; *μ*(Mo *K*α) = 0.169 mm^{−1}; *T* = 295(2) K; colorless needle; 10,657 independent measured reflections; *F*² refinement; *R*₁ = 0.0764; *wR*₂ = 0.1775.
- For more details of this synthetic method, see: Hung, W.-C.; Liao, K.-S.; Y.-H. Liu; Peng, S.-M.; Chiu, S.-H. *Org. Lett.* **2004**, *6*, 4183–4186.
- Data for 3-H-PF₆: Mp 91–93 °C; ¹H NMR (400 MHz, CD₃SOCD₃): 0.88 (t, *J* = 7 Hz, 12H), 1.29–1.38 (m, 8H), 1.51–1.58 (m, 8H), 2.47 (br, 4H), 3.32 (br, 4H), 3.58 (br, 4H), 3.75–3.89 (m, 8H), 3.90–3.95 (m, 4H), 4.24 (s, 4H), 4.62 (d, *J* = 6 Hz, 4H), 5.41–5.47 (m, 2H), 6.76 (d, *J* = 8 Hz, 4H), 6.96 (d, *J* = 8 Hz, 4H), 6.99 (d, *J* = 8 Hz, 4H), 7.12 (d, *J* = 8 Hz, 4H), 7.33 (br, 2H), 8.33 (t, *J* = 7 Hz, 1H), 8.40 (d, *J* = 7 Hz, 2H), 9.82 (br, 2H); ¹³C NMR (100 MHz, CD₃OD): 14.2, 20.1, 33.7 (*J*_{PC} = 7 Hz), 43.3, 45.5, 52.1, 67.6 (*J*_{PC} = 6 Hz), 70.8, 71.8, 75.3, 126.6, 128.4, 129.6, 129.9, 130.7, 130.9, 136.4, 141.0, 141.8, 143.0 (*J*_{PC} = 5 Hz), 150.2, 165.3; ³¹P NMR (162 MHz, CD₃OD): −138.5 (septet, *J* = 707 Hz, PF₆[−]), 15.1; HRMS (ESI): *m/z* calcd for [3-H]⁺ (C₅₉H₈₅N₆O₁₁P₂) 1115.5752; found 1115.5811.
- (a) Furusho, Y.; Matsuyama, T.; Takata, T.; Moriuchi, T.; Hirao, T. *Tetrahedron Lett.* **2004**, *45*, 9593–9597; (b) Fuller, A.-M.; Leigh, D. A.; Lusby, P. J.; Oswald, I. D. H.; Parsons, S.; Walker, D. B. *Angew. Chem., Int. Ed.* **2004**, *43*, 3914–3918.
- Crystal data for **5**: [C₃₄H₃₆N₄O₅Pd·H₂O][PF₆]; *M*_r = 705.08; monoclinic; space group *P*2₁/*n*; *a* = 12.4078(2); *b* = 9.3147(1); *c* = 28.1577(3) Å; *V* = 2194.10(7) Å³; *ρ*_{calcd} = 1.466 g cm^{−3}; *μ*(Mo *K*α) = 0.632 mm^{−1}; *T* = 295(2) K; colorless needle; 7303 independent measured reflections; *F*² refinement; *R*₁ = 0.0378; *wR*₂ = 0.0841.
- Data for **9**: Mp 247–249 °C; ¹H NMR (400 MHz, CDCl₃): 3.50–3.52 (m, 4H), 3.60–3.63 (m, 4H), 4.28 (s, 4H), 4.34 (s, 4H), 4.86 (s, 4H), 6.61 (d, *J* = 8 Hz, 4H), 6.85 (d, *J* = 8 Hz, 4H), 7.13–7.18 (m, 12H), 7.19–7.22 (m, 24H), 7.30 (d, *J* = 8 Hz, 4H), 7.42 (t, *J* = 8 Hz, 1H), 7.48–7.51 (m, 3H), 7.81 (br, 2H); ¹³C NMR (100 MHz, CDCl₃): 29.8, 50.2, 52.9, 64.5, 68.3, 70.2, 73.5, 82.9, 92.2, 117.4, 124.2, 125.7, 127.3, 127.4, 128.9, 130.8, 131.5, 135.1, 135.6, 137.8, 140.1, 140.3, 141.7, 143.9, 146.7, 152.2, 152.8, 171.0; HRMS (FAB): *m/z* calcd for [9]⁺ (C₉₀H₇₄N₆O₉Pd) 1488.4552; found 1488.4597. For more details of this synthetic method, see Ref. 15a.
- We were unable to remove the Pd(II) ion from the [2]rotaxane **9** when using catalytic hydrogenolysis (Pd/C), hydrolysis under acidic (HCl) and basic (Et₃N) conditions, or high-pressure carbon monoxide (50 atm), suggesting that relatively tight binding occurs between the components and supporting the integrity of the purported rotaxane-like molecular structure.