This article was downloaded by: [National Taiwan University]

On: 31 August 2009

Access details: Access Details: [subscription number 731897496]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK

Synthetic Communications

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title-content=t713597304>

Preparation of Functionalized Imidazolium Salts under Microwave Irradiation

Shih-Kang Fu ^a; Shih-Tzung Liu ^a

^a Department of Chemistry, National Taiwan University, Taipei, Taiwan

Online Publication Date: 01 June 2006

To cite this Article Fu, Shih-Kang and Liu, Shih-Tzung(2006)'Preparation of Functionalized Imidazolium Salts under Microwave Irradiation',Synthetic Communications,36:14,2059 — 2067

To link to this Article: DOI: 10.1080/00397910600634464

URL: <http://dx.doi.org/10.1080/00397910600634464>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Preparation of Functionalized Imidazolium Salts under Microwave Irradiation

Shih-Kang Fu and Shiuh-Tzung Liu

Department of Chemistry, National Taiwan University, Taipei, Taiwan

Abstract: A direct alkylation of a substituted imidazole to prepare the corresponding functionalized ionic liquid has been developed in excellent yields under microwave irradiation.

Keywords: Imidazolium, ionic liquid, microwave irradiation

From an environmental perspective, ionic liquids provide an alternative reaction medium for chemical reactions.^[1] Among various ionic liquids, simple alkyl substituted 1,3-dialkyl-imidazolium cations associated with weakly coordinating anions are the most popular used in biphasic catalysis because of their distinct physical–chemical properties.^[1,2] More recently, developments of the presence of appropriate functional groups as part of ionic liquid molecules to increase the immobilization of catalysts or facilitate the separation process represent a new area of interest.^[3] Thus, a few of this type of ionic liquids, such as imidazolium-based ones with amine,^[4] amido,^[5] hydroxy,^[6] carboxylate,^[7] urea,^[8] perfluoro-chains,^[9] and nitriles,^[3] are known.

Conventional heating in refluxing solvents is a typical way to prepare 1,3-dialkylimidazolium halides. Compared to the time-consuming processes of conventional methods, a microwave-assisted chemical reaction has been applied to synthesize ionic liquids.^[10,11] In the past two decades, the use of microwave ovens in chemical synthesis and analysis has grown because of the advantages in reducing reaction times, improving yield, and simplifying procedures. The preparation of functionalized ionic liquids using a solvent-free approach has not reported in literature.^[4–9]

Received in Japan March 7, 2006

Address correspondence to Shiuh-Tzung Liu, Department of Chemistry, National Taiwan University, Taipei 106, Taiwan. E-mail: stliu@ntu.edu.tw

RESULTS AND DISCUSSION

The microwave-assisted reaction was conducted in a round-bottom flask using a Discovery microwave heating apparatus with a thermal controller (CEM Corp., Matthews, NC). This apparatus allows the reaction flask to be equipped with condenser fitted into a microwave-irradiation chamber (see the experimental section). Typically, a mixture of equal molar amounts (6.5 mmol) of 1-methylimidazole and γ -bromobutyronitrile in a 20-mL flask was irradiated at 180 W at 180°C for 30 s. Upon completion of the reaction, the pale yellow, viscous oil was washed with dichloromethane (20 mL $\times$ 2) to remove the unreacted starting materials. The residue was dried under vacuum overnight, and the obtained material was analytical pure.

All reactions studied in this work are shown in Scheme 1. As a starting point, we studied the microwave-assisted coupling of 1-methylimidazole with 3-triethoxysilylpropyl chloride. Entries 1–5 of Table 1 show the results of this optimization. It appeared that the period of irradiation and reaction temperature were critical for the reaction. The best conditions for the preparation of **1** is irradiation at microwave power of 245 W for 40 s. Either a higher or a lower power of irradiation in this reaction gave decomposition or poorer yield. The scale for the reaction is irrelevant to the irradiation period employed. Because of the moisture sensitivity of the triethoxy silyl group, the reaction had to be carried out under a dry nitrogen atmosphere; otherwise the condensation leading to Si-O-Si took place, which caused a decrease in the yield of **1**. However, it can be carried out under aerobic

Scheme 1. Preparation of functionalized 1,3-disubstituted imidazolium-based ionic liquids.

Table 1. Results for the reactions of 1-methylimidazole with various alkyl halides^a

Entry	Quantity of reactant (mmol)	Product	Power of MW (W)	T (s)	Temp. (°C)	Atm.	Yield ^b	Physical property	Ref.
1	6.5	1	245	40	245	N ₂	95%	Pale yellow liquid	[12]
2	60	1	245	40	245	air	49%	Pale yellow liquid	
3	6.5	1	245	25	140	N ₂	trace	—	
4	6.5	1	185	40	145	N ₂	trace	—	
5	6.5	1	245	40	185	N ₂	93%	Pale yellow liquid	[13]
6	11.3	2	185	25	180	air	90%	Pale yellow liquid	
7	6.5	3	180	30	180	air	95%	Pale yellow liquid	[3]
8	6.5	4	100	25	175	air	92%	Lemon yellow liquid	[14]
9	6.5	5	100	35	180	air	90%	Colorless liquid	[15]
10	3.4	6	100	35	168	air	90%	Pale yellow liquid	[16]
11	4.6	7	180	30	180	air	81%	Lemon yellow liquid	
12	5.0	8	250	90	185	air	90%	Colorless liquid	
13	2.9	9	250	90	185	air	85%	Colorless liquid	
14	1.5	10	30	20	110	air	45%	White powder solid	
15	1.5	10	100	20	170	air	dec.	—	
16	1.5	10	5	20	170	air	0	—	

^aReaction conditions: an equal molar amount of 1-methylimidazole and RX in flask without using any solvent.^bIsolated yield and characterized by ¹HNMR.

conditions for other substrates. All products were characterized by ^1H NMR and elemental analysis. The optimum reaction conditions for reactions of 1-methylimidazole with various alkyl halides are summarized in Table 1.

As shown in Table 1, various alkyl halides were able to react with imidazole to give the corresponding ionic liquid in excellent yields (entries 6–13). For example, the production of **10** was achieved by adopting 30 W at 110°C , whereas the compound **8** was with 250 W at 185°C . For the acetylated glucose derivative, the reaction provided a moderate yield by using 30 W of microwave at 110°C . Attempts to improve the yield by varying the microwave powers and temperatures failed, indicating an intrinsic property of this substrate.

All functionalized imidazolium halides except **10** are in liquid phase at room temperature. Compound **10** is in a solid form, which becomes liquid at temperatures greater than $90\text{--}93^\circ\text{C}$. As for solubility, those containing —OH of polyether groups are water- and methanol-soluble materials. In fact, compound **8** is completely miscible with water. Of course, the halide anions of this series of ionic liquids also increase their solubility toward water. Similarly, the alkylation of 1-propyl-2-methylimidazole proceeded smoothly to give the corresponding ionic liquid in excellent yield (Scheme 2). The reaction conditions employed are comparable to the preparation of the related 1,3-disubstituted imidazolium-based ionic liquids (Table 2).

Reaction of 1-bromo-3-chloropropane with 1-methylimidazole under microwave irradiation yielded a mixture of **14**, **15**,^[17] and **16**^[18] (Scheme 3). It shows that the poor selectivity of nucleophilic substitution of imidazole toward alkyl bromide versus chloride under microwave-assisted conditions even with a lower power irradiation. Even under mild conditions with lower power (30 W), this reaction still provided a mixture of products. However, the reaction proceeded poorly at the temperature lower than 100°C . All these compounds were not able to be obtained in pure form because of the anion.

3-Aminopropyl chloride was also tested as a substrate for the alkylation. A mixture of an equalmolar amount of 3-aminopropyl chloride hydrochloride

Scheme 2. Preparation of functionalized 1,2,3-trisubstituted imidazolium-based ionic liquids.

Table 2. Results for preparation of 1,2,3-trisubstituted imidazolium-based ionic liquids

Product	Power (W)	T (s)	Temp. (°C)	Yield (%)	Physical property
11	235	40	185	75	Pale yellow liquid
12	180	50	180	89	Pale yellow liquid
13	105	30	180	85	Pale yellow liquid

salt, 1-methylimidazole, and triethylamine was irradiated at 170°C with the power setting of 180 W for 90 s to provide the desired ionic liquid **17** in 65% yield accompanied by the formation of azetidine Eq. (1). This outcome shows that the alkylation takes place with imidazole nitrogen superior to the amine site. However, the use of excess 3-aminopropyl chloride salt can provide **17** in a high yield.

Scheme 3. Reaction of imidazole with 3-chloro-1-bromopropane.

In summary, we have demonstrated a much faster and more efficient method for the preparation of functionalized imidazolium-based ionic liquid in high yields. Work on these functionalized ionic liquids continues and is expected to illustrate interesting new results.

EXPERIMENTAL

Nuclear magnetic resonance spectra were recorded in CDCl_3 or acetone- d_6 on either a Bruker AM-300 or Avance 400 spectrometer. Chemical shifts are given in parts per million relative to Me_4S for ^1H and ^{13}C NMR. Infrared spectra were measured on a Nicolet Magna-IR 550 spectrometer (series-II) as KBr pallets, unless otherwise noted. Microwave irradiation was carried out in a Discovery microwave heating apparatus with temperature controller (CEM Corp., Matthews, NC). The reaction temperature reported was based on this readout. Chemicals and solvents were of analytical grade and used as received unless otherwise stated. α -Bromotetraacetyl-L-glucose was prepared from the earlier reported procedures.^[19]

General Procedure

An equal molar amount of 1-methylimidazole and alkyl halides in a 20-mL flask was set up as shown in Fig. 1 and was irradiated at a certain level of power with programming the temperature for a certain period of reaction time. Upon completion of the reaction, the pale yellow, viscous oil was washed with dichloromethane ($20\text{ mL} \times 2$) to remove the unreacted starting materials and was dried under vacuum overnight. Spectroscopic data of the obtained compounds **1–6** are essentially identical to the literature reported.^[3,12–16]

Figure 1. Microwave-assisted heating apparatus.

Data

Compound 7. $\nu_{\max}$ KBr/cm⁻¹ 3164, 2952, 2866, 1726 ($\nu_{\text{C=O}}$), 1653 ($\nu_{\text{C=C}}$), 1573; ¹H NMR (300 MHz, CDCl₃): δ 9.99 (s, 1H), 7.53 (s, 1H), 7.47 (s, 1H), 4.20 (t, 2H, $J = 7.31$ Hz), 3.93 (s, 3H), 3.91 (q, 2H, $J = 7.05$ Hz), 2.16 (t, $J = 7.29$ Hz), 1.80 (m, 2H), 1.50 (m, 2H), 1.24 (m, 2H), 1.07 (t, 3H, $J = 7.18$ Hz). MS found [M-Br]⁺ 239 for C₁₃H₂₃N₂O₂; anal. calcd. for C₁₃H₂₃BrN₂O₂: C, 48.91; H, 7.26; N, 8.78. Found for C, 45.45; H, 7.15; N, 9.46.

Compound 8. $\nu_{\max}$ KBr/cm⁻¹ 3552 ($\nu_{\text{O-H}}$), 3164, 2872, 1653 ($\nu_{\text{C=C}}$), 1587, 1116; ¹H NMR (300 MHz, D₂O): δ 8.75 (s, 1H), 7.52 (s, 1H), 7.45 (s, 1H), 4.39 (t, 2H, $J = 4.6$ Hz), 3.91 (s, 3H), 3.89 (m, 2H), 3.72 (m, 6H), 3.60 (m, 2H). MS found [M-Cl]⁺ 215 for C₁₀H₁₉N₂O₃; anal. calcd. for C₁₀H₁₉ClN₂O₃: C, 47.9; H, 7.64; N, 11.19. Found: C, 47.21; H, 7.83; N, 11.37.

Compound 9. $\nu_{\max}$ KBr/cm⁻¹ 3164, 2873, 1646, 1566, 1103; ¹H NMR (300 MHz, CDCl₃): δ 9.81 (s, 1H), 7.66 (s, 1H), 7.47 (s, 1H), 3.97 (s, 3H), 3.79 (t, 2H, $J = 4.58$ Hz), 3.52 (m, 8H), 3.25 (s, 3H). MS found [M-Br]⁺ 229 for C₁₁H₂₁N₂O₃; anal. calcd. for C₁₁H₂₁BrN₂O₃: C, 42.73; H, 6.85; N, 9.06. Found: C, 42.59; H, 7.19; N, 8.98.

Compound 10. Mp: 90–93°C; $\nu_{\max}$ KBr/cm⁻¹ 3171, 2939, 2879, 1726 ($\nu_{\text{C=O}}$), 1653, 1587, 1089; ¹H NMR (300 MHz, CDCl₃): δ 10.05 (s, 1H), 7.38 (s, 1H), 7.31 (s, 1H), 5.15 (t, 1H, $J = 9.6$ Hz), 5.02 (t, 1H, $J = 9.9$ Hz), 4.85 (t, 1H, $J = 8.0$ Hz), 4.50 (d, 1H, $J = 8.0$ Hz), 4.39 (m, 2H), 4.20 (m, 1H), 4.06 (s, 3H), 3.71 (m, 2H), 3.62 (m, 2H), 2.20 (m, 2H), 2.05 (s, 1H), 2.02 (s, 1H), 2.00 (s, 1H), 1.95 (s, 1H). MS found [M-Br]⁺ 472 for C₂₀H₂₉N₂O₁₀; anal. calcd. for C₂₀H₂₉BrN₂O₁₀: C, 44.70; H, 5.44; N, 5.21. Found: C, 44.26; H, 5.45; N, 4.99.

Compound 11. $\nu_{\max}$ KBr/cm⁻¹ 3144, 2980, 2933, 2849, 1640, 1586, 1076; ¹H NMR (300 MHz, CDCl₃): δ 7.66 (s, 1H), 7.51 (s, 1H), 4.27 (t, 2H, $J = 7.3$ Hz), 4.08 (t, 6H, $J = 7.3$ Hz), 3.92 (q, 2H, $J = 7.2$ Hz), 2.65 (s, 3H), 2.34 (t, 2H, $J = 7.2$ Hz), 1.97 (m, 2H), 1.72 (m, 2H), 0.82 (t, 9H, $J = 7.3$ Hz). MS found [M-Cl]⁺ 287 for C₁₃H₂₇N₂O₃Si; anal. calcd. for C₁₃H₂₇ClN₂O₃Si: C, 48.35; H, 8.43; N, 8.68. Found: C, 47.21; H, 7.83; N, 11.37.

Compound 12. $\nu_{\max}$ KBr/cm⁻¹ 3403 ($\nu_{\text{O-H}}$), 3144, 2952, 2886, 1653, 1593, 1129; ¹H NMR (300 MHz, CDCl₃): δ 7.81 (s, 1H), 7.49 (s, 1H), 4.43 (t, 2H, $J = 4.5$ Hz), 4.10 (t, 2H, $J = 7.2$ Hz), 3.81 (t, 2H, $J = 4.5$ Hz), 3.52 (m, 8H), 2.65 (s, 3H), 1.75 (m, 2H), 0.88 (t, 3H, $J = 5.2$ Hz). MS found [M-Cl]⁺ 257 for C₁₃H₂₅ClN₂O₃; anal. calcd. for C₁₃H₂₅ClN₂O₃: C, 53.33; H, 8.61; N, 9.57. Found: C, 53.18; H, 8.98; N, 9.44.

Compound 13. $\nu_{\max}$ KBr/cm⁻¹ 3139, 2974, 2883, 1727, 1640, 1587; ¹H NMR (300 MHz, CDCl₃): δ 7.65 (s, 1H), 7.54 (s, 1H), 4.19 (t, 2H, J = 11.2 Hz), 3.43 (s, 9H), 2.72 (s, 3H), 1.79 (m, 2H), 0.84 (t, 3H, J = 18.4 Hz), 0.54 (t, 3H, J = 12.6 Hz). MS found [M-Br]⁺ 239 for C₁₃H₂₃BrN₂O₂; anal. calcd. for C₁₃H₂₃BrN₂O₂: C, 48.91; H, 7.26; N, 8.78. Found for C, 48.96; H, 7.93; N, 8.96.

Compound 17. $\nu_{\max}$ KBr/cm⁻¹ 3455 ($\nu_{\text{N-H}}$), 3157, 2963, 2753, 1634, 1579; ¹H NMR (300 MHz, D₂O): δ 8.65 (s, 1H), 7.39 (s, 1H), 7.32 (s, 1H), 4.19 (t, 2H, J = 7.3 Hz), 3.75 (s, 3H), 2.93 (m, 2H), 2.16 (m, 2H). MS found [M-Cl]⁺ 140 for C₇H₁₄N₃; anal. calcd. for C₇H₁₄ClN₃: C, 47.86; H, 8.03. Found for C, 47.66; H, 7.83.

ACKNOWLEDGMENT

This research was supported by the National Science Council (NSC93-2113-M002-004), Taiwan, China.

REFERENCES

1. Recent reviews: (a) Wilkes, J. S. Properties of ionic liquid solvents for catalysis. *J. Mol. Catal. A, Chem* **2004**, *214*, 11–17; (b) Dzyuba, S. V.; Bartsch, R. A. Recent advances in applications of room-temperature ionic liquid/supercritical CO₂ systems. *Angew. Chem. Int. Ed.* **2003**, *42*, 148–150; (c) Dupont, J.; de Souza, R. F.; Suarez, P. A. Z. Ionic liquid (molten salt) phase organometallic catalysis. *Chem. Rev.* **2002**, *102*, 3667–3692; (d) Handy, S. T. Greener solvents: Room temperature ionic liquids from biorenewable sources. *Chem.-Eur. J.* **2003**, *9*, 2938–2944.
2. Suarez, P. A. Z.; Dullius, J. E. L.; Einloft, S.; de Souza, R. F.; Dupont, J. Synthesis and physical-chemical properties of ionic liquids based on 1-n-butyl-3-methylimidazolium cation. *J. Chim. Phys.* **1998**, *95*, 1626–1639.
3. Zhao, D.; Fei, Z.; Scopelliti, R.; Dyson, P. J. Synthesis and characterization of ionic liquids incorporating the nitrile functionality. *Inorg. Chem.* **2004**, *43*, 2197–2205.
4. Herrmann, W. A.; Köcher, C.; Goossen, L. J.; Artus, G. R. J. Heterocyclic carbenes: A high-yielding synthesis of novel, functionalized N-heterocyclic carbenes in liquid ammonia. *Chem.-Eur. J.* **1996**, *2*, 1627–1636.
5. Lee, K.-M.; Lee, Y.-T.; Lin, I. J. B. Supramolecular liquid crystals of amide functionalized imidazolium salts. *J. Mater. Chem.* **2003**, *13*, 1079–1084.
6. Branco, L. C.; Rosa, J. N.; Moura Ramos, J. J.; Alfons, C. A. M. Preparation and characterization of new room temperature ionic liquids. *Chem.-Eur. J.* **2002**, *8*, 3671–3677.
7. (a) Cole, A. C.; Jensen, J. L.; Ntai, I.; Tran, K. L. T.; Weaver, K. J.; Forbes, D. C.; Davis, J. H., Jr. Novel Brønsted acidic ionic liquids and their use as dual solvent-catalysts. *J. Am. Chem. Soc.* **2002**, *124*, 5962–5963; (b) Holbrey, J. D.; Reichert, W. M.; Tkatchenko, I.; Bouajila, E.; Walter, O.; Tommasi, I.

- Roger, R. D. 1,3-Dimethylimidazolium-2-carboxylate: The unexpected synthesis of an ionic liquid precursor and carbene-CO₂ adduct. *Chem. Commun.* **2003**, 28–29.
8. Visser, A. E.; Swatloski, R. P.; Reichert, W. M.; Mayton, R.; Sheff, S.; Wierzbicki, A.; Davis, J. H., Jr.; Rogers, R. D. Task-specific ionic liquids for the extraction of metal ions from aqueous solutions. *Chem. Commun.* **2001**, 135–136.
9. Merrigan, T. L.; Bates, E. D.; Dorman, S. C.; Davis, J. H., Jr. New fluororous ionic liquids function as surfactants in conventional room-temperature ionic liquids. *Chem. Commun.* **2000**, 2051–2502.
10. (a) Varma, R. S.; Nambodiri, V. V. Solvent-free preparation of ionic liquids using a household microwave oven. *Pure Appl. Chem.* **2001**, *73*, 1309–1313; (b) Nambodiri, V. V.; Varma, R. S. Microwave-assisted preparation of dialkylimidazolium tetrachloroaluminates and their use as catalysts in the solvent-free tetrahydropyranlation of alcohols and phenols. *Chem. Commun.* **2002**, 342–343.
11. (a) Thanh, G. V.; Pegot, B.; Loupy, A. Solvent-free microwave-assisted preparation of chiral ionic liquids from (-)-N-methylephedrine. *Eur. J. Org. Chem.* **2004**, 1112–1116; (b) Ranu, B. C.; Jana, R.; Dey, S. S. An efficient and green synthesis of 2-arylbenzothiazoles in an ionic liquid, [pmIm]Br under microwave irradiation. *Chem. Lett.* **2004**, *33*, 274–275; (c) Peng, Y.; Song, G. Microwave-assisted clean synthesis of 6-aryl-2,4-diamino-1,3,5-triazines in [bmim][PF₆]. *Tetrahedron Lett.* **2004**, *45*, 5313–5316.
12. Chi, Y. S.; Lee, J. K.; Lee, S.-G.; Choi, I. S. Control of wettability by anion exchange on Si/SiO₂ surfaces. *Langmuir* **2004**, *20*, 3024–3027.
13. Ricciardi, F.; Romanchick, W. A.; Joullie, M. M. 1,3-Dialkylimidazolium salts as latent catalysts in the curing epoxy-resins. *J. Polym. Sci., Polym. Lett.* **1983**, *21*, 633–638.
14. Walker, A. J.; Bruce, N. C. Combined biological and chemical catalysis in the preparation of oxycodone. *Tetrahedron* **2004**, *60*, 561–568.
15. Dubreuil, J. F.; Bazureau, J. P. Rate accelerations of 1,3-dipolar cycloaddition reactions in ionic liquids. *Tetrahedron Lett.* **2000**, 7351–7355.
16. Horvath, A. Hofmann-type elimination in the efficient N-alkylation of azoles-imidazole and benzimidazole. *Synthesis* **1994**, 102–106.
17. Zhang, D.; Chen, J.; Liang, Y.; Zhou, H. Facile synthesis of novel ionic liquids containing dithiocarbamate. *Synth. Commun.* **2005**, *35*, 521–526.
18. Poyatos, M.; Mas-Marza, E.; Sanau, M.; Peris, E. Synthesis and reactivity of new chelate-N-heterocyclic biscarbene complexes of ruthenium. *Inorg. Chem.* **2004**, *43*, 1793–1798.
19. Potter, A. L.; Sowden, J. C.; Hassid, W.; Doudoroff, M. α -L-Glucose-1-phosphate. *J. Am. Chem. Soc.* **1948**, *70*, 1751–1752.