LETTERS AND COMMENTS

Documentation forms for patients with asthma: an evaluation

TO THE EDITOR: The focus of pharmacy practice is moving from being centered on dispensing to being focused on the patient. Professional organizations are promoting patient-centered care as routine, for which systematic and consistent documentation is essential. We report on an evaluation of the use of various documentation methods by community pharmacists managing patients with asthma, with a view to developing documentation in Portuguese pharmacy practice.

Methods. The study was based in 5 community pharmacies during February and March 2002. Patients using the pharmacy who were prescribed asthma medication were invited to participate in the study. Patients agreed to participate following receipt of a leaflet explaining the study and signed a consent form. The pharmacists' interventions in managing these patients were documented using various data collection forms. These included a questionnaire to predict respiratory conditions in adults (IUALTD)¹; a form to document patient data, the drugs prescribed, and drug-related problem (DRP) classification²; the Mini-Asthma Quality of Life Questionnaire³; a checklist to assess patients' inhaler techniques⁴; a questionnaire to assess the patients' knowledge of their drugs and condition⁵; and a patient information leaflet developed by the research team. Before the study, pharmacists attended a training session to familiarize themselves with the documentation. The pharmacists' opinions of the use and utility of the documentation were evaluated using a 5-point scale. Descriptive and inferential statistics (SPSS for Windows, release 10) were used to analyze the collected data.

Results. The participating pharmacists developed opinions about the documentation tools after using them over the 2-month study period on 24 patients. Every tool was considered generally useful (Table 1). However, the DRP classification was useful but not very simple to use and generated considerable workload. The lowest scores were obtained for the diagnostic tool and the knowledge and quality of life questionnaire ($\bar{x} = 3.0$), both of which are often used in research studies and therefore may need to be reviewed if research is to be fully implemented in prac-

tice. While the inhaler technique checklist was considered useful and simple, it created extra workload. Three of the piloted tools were regarded as able to be incorporated in daily practice. Pharmacists ranked the patient information leaflets highest (4.9), considering them to be very useful and simple while generating no extra workload. The use of peak flow meters was also ranked very positively (4.3), viewed as very useful and quite simple to use, even though they generated some extra workload. The patient profile forms were ranked quite positively (3.9); they were useful and simple, while creating some workload.

The limitations of this study include the small sample and the convenience sampling process used. The study may not be widely generalizable; however, it is important to report on such feasibility studies to help with understanding how best to target such implementation strategies. It seems that most of the methods of documentation are considered usable in daily practice and could be made part of the documentation processes being implemented in professional activities in Portugal.

Funding for the study was provided by Faculdade de Farmácia da Universidade de Lisboa and Associação Nacional das Farmácias; GlaxoSmithKline provided the IUALTD and the peak flow meters. We thank patients and pharmacists who actively participated in the study.

Filipa A Costa MPharm MRPharmS MSc

Health Researcher
CEFAR, Associação Nacional das Farmácias
ANF, R. Marechal Saldanha, 1; 1249-069
Lisbon, Portugal
fax 351 213400674
filipa.costa@anf.pt

JW Foppe van Mil PharmD PhD

Pharmacy Practice Consultant
van Mil Consultancy/Quality Institute for
Pharmaceutical Care
Zuidlaren, Netherlands

Catherine A Duggan BPharm (Hons) PhD MRPharmS
Director
Academic Department of Pharmacy
Barts and The London NHS Trust
London, United Kingdom

Pedro L Mata MD

Medical Doctor, Imunoallergologist
British Hospital
Lisboa, Portugal

Margarida Caramona PharmD PhD
Professor of Pharmacology
Laboratório de Farmacologia da Faculdade de
Farmácia da Universidade de Coimbra
Coimbra, Portugal

Published Online, 26 Oct 2004, www.theannals.com
DOI 10.1345/aph.1E059

REFERENCES

1. Bai J, Peat JK, Berry G, Marks GB, Woolcock AJ. Questionnaire items that predict asthma and other respiratory conditions in adults. *Chest* 1998;114:1343-8.

Table 1. Ranking of Tested Tools

Tool	Utility ^a	Simplicity ^b	Workload ^c	Overall ^d
IUALTD	4.0	3.7	2.6	3.1
PFM	4.7	4.4	4.4	4.3
Mini-AQLQ	3.9	3.6	2.9	2.9
Patient profile	4.3	3.7	2.7	3.9
DRP classification	4.2	2.7	2.2	3.0
Inhaler technique checklist	4.2	3.8	3.5	3.3
Knowledge questionnaire	3.7	3.0	3.2	3.0
Patient information leaflets	4.9	4.7	4.9	4.9

AQLQ = Asthma Quality of Life Questionnaire; DRP = drug-related problem; IUALTD = International Union Against Tuberculosis and Lung Disease; PFM = peak flow meter.

^a5. Very useful; 4. useful; 3. neither useful nor useless; 2. little useful; 1. useless.

^b5. Very simple; 4. simple; 3. neither simple nor complicated; 2. complicated; 1. very complicated.

^c5. Does not generate any workload; 4. does not generate a lot of workload; 3. does not increase or diminish workload; 2. generates some workload; 1. generates a lot of workload.

^dOverall opinion had the following introductory statement: "I'd like to have this material available for daily use in the pharmacy" (5. I strongly agree; 4. I agree; 3. I don't agree or disagree; 2. I disagree; 1. I strongly disagree).

- van Mil JWF, Winterstein A. Documenting system for drug related problems. Preliminary Validation Report v1.2. Pharmaceutical Care Network Europe 2001. www.pcne.org (accessed 2003 Feb 19).
- Juniper EF, Guyatt GH, Cox FM, Ferrie PJ, King DR. Development and validation of the Mini-Asthma Quality of Life Questionnaire. *Eur Respir J* 1999;14:32-8.
- Pharmacy-based asthma services. Protocol and guidelines 1998. Geneva: World Health Organization, 1998.
- Questionnaire de connaissances sur l'asthme et son traitement. Santé Education Recherche Action. www.respir.com (accessed 2002 Jan 14).

Fluorouracil-induced neurotoxicity presenting with generalized tonic-clonic seizure

TO THE EDITOR: Fluorouracil is a chemotherapeutic agent commonly used in a number of solid malignancies, including colorectal cancer. Fluorouracil is a pyrimidine antimetabolite that blocks DNA synthesis and inhibits RNA processing and function. Fluorouracil readily penetrates the blood-brain barrier. Fluorouracil-induced neurotoxicity is rare. Neurotoxicity occurs in <5% of patients receiving high doses or lengthy, continuous infusions.¹

Case Report. A 56-year-old man with a diagnosis of sigmoid adenocarcinoma with multiple liver metastases began therapy following surgery consisting of oxaliplatin 85 mg/m² on day 1, folinic acid 500 mg/m² on days 1 and 2, and continuous infusion of fluorouracil 1600 mg/m² for 48 hours, repeated every 15 days. Intravenous granisetron 3 mg was also administered during the first day of the chemotherapy cycle. He developed generalized tonic-clonic seizures during the second day of treatment; fluorouracil treatment was stopped and phenytoin 15 mg/kg was administered. No further seizures occurred after he commenced regular oral phenytoin.

The patient had no underlying medical conditions that might have induced this seizure and had no past history of seizure. On examination after the seizures, there were no neurologic abnormalities. At that time, complete blood cell count, blood glucose, serum electrolytes, renal and liver function studies, and calcium concentrations were all within normal limits. Magnetic resonance imaging of the brain with contrast showed no evidence of stroke or metastases. The electroencephalogram was normal. No adverse effects were observed during chemotherapy cycles 2-5. Phenytoin concentrations were monitored between the cycles and were within normal range. The patient discontinued phenytoin after the fifth chemotherapy cycle. However, tonic-clonic seizures recurred while the patient was receiving the seventh cycle of treatment.

Discussion. Oxaliplatin, granisetron, and folinic acid may also cause neurotoxicity. Because oxaliplatin and granisetron were administered on the first day of the cycle and the event happened during the second day of treatment, these drugs are unlikely to have been causative agents. Most cases of neurotoxicity related to folinic acid have been reported when it was given concomitantly with fluorouracil. Although there was no concomitant use of phenytoin and fluorouracil during these 2 events, phenytoin toxicity related to phenytoin-fluorouracil interaction has been reported in the literature.² Fluorouracil-induced neurotoxicity includes acute cerebellar syndrome, somnolence, and confusion.¹ However, generalized tonic-clonic seizures alone as an initial symptom of fluorouracil-induced neurotoxicity has not been reported. Based on the Naranjo probability scale, fluorouracil was a possible cause of the seizure.³

The pathogenesis of neurotoxicity associated with fluorouracil administration is not well recognized. Postulated mechanisms for the pathogenesis include Krebs cycle blockade by fluoroacetate and fluorocitrate, thiamine deficiency,⁴ and inherited deficiency of the enzyme dihydropyrimidine dehydrogenase.⁵ Acute toxicity is dose related and generally reversible. Neurotoxic effects are dose and schedule dependent and are reversible with drug withdrawal and dose reduction. Since fluorouracil is the most frequently used chemotherapy for the treatment of colorectal cancer, fluorouracil-induced neurotoxicity should be considered as an important differential diagnosis in cancer patients who develop neurological abnormality while receiving fluorouracil-based chemotherapy.

High-dose fluorouracil/folinic acid infusion therapy has recently become a popular regimen for various cancers. It is necessary that both oncologists and neurologists be fully aware of this unusual complication.

Emin Tamer Elkiran MD

Fellow
Department of Medical Oncology
Hacettepe University Faculty of Medicine
Ankara, Turkey

Kadri Altundag MD

Associate Professor
Department of Medical Oncology
Hacettepe University Faculty of Medicine
Ankara
8181 Fannin St., No: 728
Houston, Texas 77054-2916
fax 713/794-4385
drkadri@usa.net

Yavuz Beyazit MD

Resident
Department of Internal Medicine
Hacettepe University Faculty of Medicine

Nilufer Guler MD

Professor
Department of Medical Oncology
Hacettepe University Faculty of Medicine

Ayşe Kars MD

Professor
Department of Medical Oncology
Hacettepe University Faculty of Medicine

Published Online, 19 Oct 2004, www.theannals.com
DOI 10.1345/aph.1E070

REFERENCES

- Bofill JS, Chaves M, Moreno JA. Fluorouracil-induced aphasia (letter). *Ann Pharmacother* 2000;34:955. DOI 10.1345/aph.19265
- Gilbar PJ, Brodribb TR. Phenytoin and fluorouracil interaction. *Ann Pharmacother* 2001;35:1367-70. DOI 10.1345/aph.1A051
- Naranjo CA, Busto U, Sellers EM, Sandor P, Ruiz I, Roberts EA, et al. A method for estimating the probability of adverse drug reactions. *Clin Pharmacol Ther* 1981;30:239-45.
- Aksoy M, Basu T, Brint J, Dickerson JWT. Thiamine status of patients treated with drug combinations containing 5-fluorouracil. *Eur J Cancer* 1980;16:1041-5.
- Celik I, Kars A, Guc D, Tekuzman G, Ruacan S. Dihydropyrimidine dehydrogenase enzyme deficiency: clinical and genetic assessment of prevalence in Turkish cancer patients. *Cancer Invest* 2002;20:333-9.

Updated clinical risk factor analysis of suspected hypersensitivity reactions to abacavir

TO THE EDITOR: Hypersensitivity to abacavir is a well-characterized clinical syndrome typified by the appearance of symptoms indicating multi-organ/body system involvement occurring in approximately 5% of abacavir-treated patients.¹ The mechanism of the abacavir hypersensitivity reaction (HSR) has not been proven, and clinical risk factors have not been consistently identified; therefore, the identification of patients at risk remains an empiric process.

A previous retrospective analysis to identify clinical risk factors for abacavir hypersensitivity was conducted on 5332 abacavir-treated subjects, including 197 cases of suspected abacavir HSR.² In this analysis, the risk of developing HSR was lower among subjects of African descent and antiretroviral therapy-experienced subjects.

Since then, several changes in our approach to diagnosis, reporting, and management of abacavir HSR have occurred. GlaxoSmithKline implemented a standardized abacavir HSR case report form into all new protocols to collect more comprehensive information about the syndrome. Investigators were also instructed to report all suspected abacavir HSR cases as serious adverse events regardless of severity. Finally, the abacavir product label, all educational materials, and protocols were updated: respiratory symptoms were made more prominent in the definition of the syndrome and new guidance instructed prescribers to discontinue abacavir as soon as an abacavir HSR was suspected, regardless of the presence of other potential diagnoses.

To assess the potential impact of such changes and identify clinical risk factors for developing suspected abacavir HSR, GlaxoSmithKline conducted an updated retrospective analysis comprised of 8038 abacavir-treated subjects with a 5% (403 suspected HSR/8038 subjects) incidence rate of

suspected HSR (Figure 1). The most notable difference in results between these 2 analyses was the incidence of abacavir HSR reporting. Before the changes were instituted, the incidence was 3.6% compared with 7.6% afterward ($p < 0.001$), suggesting an increased awareness and potential misclassification of the reaction, as shown in double-blind studies.³ In addition, in this updated analysis, female gender was a significant prognostic factor for developing a suspected abacavir HSR (OR = 1.42).

In both analyses, subjects of African descent and patients with more advanced disease had decreased incidences of suspected abacavir HSR. In the updated analysis,⁴ subjects of African descent had 0.5 times lower odds compared with other ethnic origins; one explanation may be genetic factors that modulate the metabolism of or immune responses to abacavir or its metabolites. The odds of a subject who was Centers for Disease Control and Prevention (CDC) Class C at the time of abacavir initiation developing a suspected HSR was 0.74. CDC Class C subjects are likely to be antiretroviral therapy-experienced and therefore potentially more tolerant of adverse events.

Importantly, in this updated analysis, the strongest predictor for a reported abacavir HSR was the use of the abacavir HSR case report form module (Figure 2), likely a surrogate marker for the more conservative approach to diagnosis, management, and reporting of abacavir HSR. However, despite these statistically significant findings, no changes in the current diagnosis and management of abacavir HSR are recommended because the differences between subgroups were small and not likely to influence prescribing decisions.

The results of this study were presented in part in Abstract H-2013 at the 43rd Interscience Conference on Antimicrobial Agents and Chemotherapy, Chicago, IL, September 14–17, 2003.

Figure 1. Percentage of subjects reported as having abacavir hypersensitivity reactions (HSRs) in 34 GlaxoSmithKline protocols with and without the abacavir hypersensitivity case report form (CRF) module. Protocols marked with ◊ used the HSR CRF module.

Figure 2. Odds ratio estimates and 95% confidence intervals of significant ($p < 0.05$) stepwise selected predictors in a multivariable model using 7818 reports. 95% CIs were use of abacavir hypersensitivity case report form (HSR CRF) module (1.78 to 2.76), African ethnicity (0.40 to 0.69), male gender (0.55 to 0.90), and Centers for Disease Control and Prevention (CDC) Class C (0.58 to 0.94).

Amy G Cutrell MS

Associate Director
Statistics and Programming
GlaxoSmithKline
Research Triangle Park, North Carolina

Jaime E Hernandez MD

Senior Director & Medical Monitor
Infectious Diseases Medicine Development Center—HIV
GlaxoSmithKline
Five Moore Drive
Research Triangle Park, North Carolina 27709-3398
fax 919/315-6029
jaime.e.hernandez@gsk.com

Julie W Fleming PharmD

Lead Clinical Development Scientist
GlaxoSmithKline

Mark T Edwards MS

Statistician
Tomorrow Associates, Inc.
Cary, North Carolina

Martha Anne Moore BSPharm

Director
Regulatory Affairs
GlaxoSmithKline

Cindy H Brothers MSPH

Associate Director
GlaxoSmithKline

Trevor R Scott PhD

Group Director
Infectious Diseases Medicine Development Center—HIV
GlaxoSmithKline

Published Online, 9 Nov 2004, www.theannals.com
DOI 10.1345/aph.1E202

REFERENCES

- Hetherington S, McGuirk S, Powell G, Cutrell A, Naderer O, Spreen B, et al. Hypersensitivity reactions during therapy with the nucleoside reverse transcriptase inhibitor abacavir. *Clin Ther* 2001;23:1603-14.
- Symonds W, Cutrell A, Edwards M, Steel H, Spreen B, Powell W, et al. Risk factor analysis of hypersensitivity reactions to abacavir. *Clin Ther* 2002;24:565-73.
- Hernandez J, Cutrell A, Bonny T, Castillo S, Brothers C, Hee J, et al. Diagnosis of abacavir hypersensitivity reactions among patients not receiving abacavir in two blinded studies (abstract 134). Presented at: 5th International Workshop on Adverse Drug Reactions and Lipodystrophy in HIV, Paris, July 8–11, 2003.
- Hetherington S, Hughes AR, Mosteller M, Shortino D, Backer KL, Spreen W, et al. Genetic variations in HLA-B region and hypersensitivity reactions to abacavir (letter). *Lancet* 2002;359:1121-2.

Association of aspirin with eosinophilia in peripheral blood

TO THE EDITOR: Aspirin is widely used in the treatment of pain, inflammation, and fever, as well as for prophylaxis of myocardial infarction and stroke. Its association with asymptomatic peripheral eosinophilia has not been previously reported.

Case Report. A 75-year-old man was diagnosed with Parkinson's disease in July 1997; aspirin 100 mg daily was started one month later since a computed tomographic scan of the brain showed lacunar stroke. No baseline eosinophil count was obtained at that time. In March 1999, he was admitted because of altered mental status due to the adverse effects of trihexyphenidyl for Parkinson's disease. After this medicine was discontinued, the patient did well without any sequelae. On that admission, his lab-

Table 1. Association of AEC and Aspirin Use

Date	AEC (cells/mm ³)	Event
8/18/1997	NA	start aspirin
3/14/1999	108	admission
10/22/2002	4734	health check-up
11/06/2002	NA	discontinue aspirin
11/19/2002	900	follow-up
1/10/2003	365	follow-up

AEC = absolute eosinophil count; NA = not available.

oratory tests revealed a leukocyte count of 5.40×10^3 cells/mm³ with 2% eosinophils (absolute eosinophil count [AEC] 108 cells/mm³). He had a health screening in October 2002, and laboratory tests showed a leukocyte count of 7.89×10^3 cells/mm³ with 60% eosinophils (AEC 4734 cells/mm³) (Table 1). Besides aspirin, the medications the man was taking included levodopa 100 mg and pergolide mesylate 0.25 mg 3 times daily, docusate sodium 100 mg and the laxative Sterculia 7 g once daily, and finasteride 5 mg at bedtime.

The studies for eosinophilia did not reveal evidence of bronchial asthma, allergic rhinitis, eczema, urticaria, collagen vascular diseases, or helminthic infection. The patient underwent chest X-ray, abdominal ultrasound, esophagogastroduodenoscopic examinations, and a whole body positron emission tomographic scan. All studies for possible malignancy were unrevealing.

A search of databases including Micromedex, Iowa Drug Information System, MEDLINE, and EMBASE indicated that aspirin and levodopa have been linked to drug-induced eosinophilia.¹⁻⁶ Levodopa was critical for the control of our patient's Parkinson's disease. Aspirin was therefore withheld first. The AEC reduced dramatically at that time to 900 cells/mm³ in 2 weeks and 365 cells/mm³ in 10 weeks. During this period, all other medications were continued. The Naranjo probability scale rated the adverse drug reaction at the level of highly possible.⁷

Discussion. Previous studies have shown a relationship between aspirin use and tissue infiltration of eosinophils in patients with bronchial asthma, nasal polyps, or hypereosinophilic syndrome.¹⁻⁴ However, the evidence showing an association between aspirin use and peripheral eosinophilia has been limited. Zeitz² described peripheral eosinophilia as one of the clinical features in aspirin-induced Samter's syndrome, and its onset may be slow. However, our patient with peripheral eosinophilia did not have other associated clinical features. In the literature, the association of eosinophilia with hypersensitivity vasculitis secondary to levodopa is also listed.⁵ This association seems to be weak in our case because the AEC dropped after withholding aspirin, while levodopa was continued. Previous studies have shown that eosinophilia is related with the overexpression of interleukin-5.^{4,6} However, the relevance between aspirin-induced eosinophilia and a systemic level of interleukin-5 remains to be determined. Based on our case, eosinophilia in the absence of clinical symptoms is a highly possible adverse event caused by aspirin, and monitoring of the patient's complete blood cell count needs to be considered.

Tsun-I Cheng MD MPH

Attending Physician
Department of Medicine
Sun Yat-Sen Cancer Center
125 Lih-Der Road
Taipei, Taiwan 112
fax 011886-2-2897-2233
ticheng@mail.kfcc.org.tw

Tsun-Jen Cheng MD ScD

Professor
Institute of Occupational Medicine and Industrial Hygiene
College of Public Health
National Taiwan University
Taipei

Shao C Chiang PharmD

Clinical Coordinator
Department of Pharmacy
Sun Yat-Sen Cancer Center

We thank Dr. Steven M Holland for his valuable discussion.

Published Online, 9 Nov 2004, www.theannals.com
DOI 10.1345/aph.1E156

REFERENCES

- Holland SM, Gallin JI. Disorders of granulocytes and monocytes. In: Braunwald E, Fauci AS, Kasper DL, Hauser SL, Longo DL, Jameson JL, eds. *Harrison's principles of internal medicine*. 15th ed. New York: McGraw-Hill, 2001:366-74.
- Zeitz HJ. Bronchial asthma, nasal polyp, and aspirin sensitivity: Samter's syndrome. *Clin Chest Med* 1988;9:567-76.
- Sanak M, Szczeklik A. Genetics of aspirin induced asthma. *Thorax* 2000;55(suppl 2):S45-7.
- Schwartz RS. The hypereosinophilic syndrome and the biology of cancer. *N Engl J Med* 2003;348:1199-214.
- Levodopa. In: Klasco RK, ed. *DRUGDEX System*. Greenwood Village, CO: Thomson MICROMEDEX (edition expired Sept 2004).
- Simon HU, Plotz SG, Dummer R, Blaser K. Abnormal clones of T cells producing interleukin-5 in idiopathic eosinophilia. *N Engl J Med* 1999;341:112-20.
- Naranjo CA, Busto U, Sellers EM, Sandor P, Ruiz I, Roberts EA, et al. A method for estimating the probability of adverse drug reactions. *Clin Pharmacol Ther* 1981;30:239-45.

Generalized exfoliative dermatitis induced by interferon alfa

TO THE EDITOR: Interferon (IFN) was the first drug to demonstrate good efficacy in the treatment of chronic hepatitis C virus (HCV) infection.^{1,2} Combination therapy with pegylated IFN alfa-2b plus ribavirin is associated with a response rate of 50% among patients infected with HCV genotype 1. Systemic cutaneous manifestations have occasionally been reported, mostly involving small and limited lesions.^{3,4}

We report a case of generalized exfoliative dermatitis and intense pruritus during therapy with pegylated IFN alfa-2b in a patient with chronic HCV infection.

Case Report. A 41-year-old man with chronic HCV infection presented with an increase in serum alanine aminotransferase and serum aspartate aminotransferase values (Table 1). There was no history of drug use or abuse. Antiviral therapy was initiated with pegylated IFN alfa-2b plus ribavirin. After 3 months, normalization of aminotransferase levels and serum HCV-RNA occurred (Table 1), but with development of urticaria and cutaneous desquamation of all of the patient's body except his face. He was taking no other medication at that time. Blood chemical evaluation did not reveal serologic evidence of autoimmune disorders. Antimicrobial antibody, anti-nuclear antibody, anti-smooth-muscle antibodies, thyroid peroxidase antibodies, and anti-double-stranded DNA were negative. Tests for total and differential circulating immunocomplexes were negative.

Ribavirin was promptly stopped and corticosteroid topical therapy for these lesions was started, without any significant improvement. Therefore, 2 weeks later, pegylated IFN alfa-2b therapy was discontinued. One week later, a new clinical evaluation showed significant decrease in cutaneous lesions. Pegylated IFN alfa-2b rechallenge was performed in the presence of clinical specialists and confirmed the development of systemic cutaneous lesions with pruritus. The Naranjo probability scale indicated a highly probable relationship between the skin manifestations and pegylated IFN alfa-2b therapy in our patient.⁵

Table 1. Laboratory Findings After the Start of Pegylated IFN alfa-2b Therapy

Variable	Timepoint (after the start of therapy)		
	0	1 wk	3 mo
AST (U/L)	189	150	29
ALT (U/L)	442	348	31
γ-GT (U/L)	215	185	22
HCV-RNA (IU/mL)	600 000		<50

ALT = alanine aminotransferase; AST = aspartate aminotransferase; γ-GT = γ-glutamyl transpeptidase; HCV = hepatitis C virus; IFN = interferon.

Assistant

Clinical Pharmacology and Pharmacovigilance Unit
Department of Experimental and Clinical Medicine
Faculty of Medicine and Surgery
University Magna Graecia of Catanzaro
Regional Pharmacovigilance Center
Mater Domini University Hospital
Via T Campanella 115, 88100
Catanzaro, Italy
fax 390961774424
luca_gallelli@hotmail.com

Maria Ferraro MD

Clinical Specialist
Internal Medicine Operative Unit
San Giovanni in Fiore Hospital
Cosenza, Italy

Gaetano F Mauro MD

Clinical Specialist
Internal Medicine Operative Unit
San Giovanni in Fiore Hospital

Giovambattista De Sarro MD FP

Full Professor
Clinical Pharmacology and Pharmacovigilance Unit
Department of Experimental and Clinical Medicine
Faculty of Medicine and Surgery
University Magna Graecia of Catanzaro
Regional Pharmacovigilance Center
Mater Domini University Hospital

Published Online, 2 Nov 2004, www.theannals.com
DOI 10.1345/aph.1E326

REFERENCES

- Poynard T, Leroy V, Mathurin P, Cohard M, Opolon P, Zarski JP. Treatment of chronic hepatitis C by interferon for longer duration than six months. *Dig Dis Sci* 1996;41(12 suppl):99S-102S.
- Carithers RL Jr, Emerson SS. Therapy of hepatitis C: meta-analysis of interferon alfa-2b trials. *Hepatology* 1997;26(3 suppl 1):83S-8S.
- Cnudde F, Gharakhanian S, Luboinski J, Dry J, Rozenbaum W. Cutaneous local necrosis following interferon injections (letter). *Int J Dermatol* 1991;30:147.
- Gallelli L, Guadagnino V, Caroleo B, Marigliano N, De Sarro G. Cutaneous ulceration induced by interferon alfa (letter). *Ann Pharmacother* 2004;38:173-4. DOI 10.1345/aph.1D198
- Naranjo CA, Busto U, Sellers EM, Sandor P, Ruiz I, Roberts EA, et al. A method for estimating the probability of adverse drug reactions. *Clin Pharmacol Ther* 1981;30:239-45.

Data mining, drug safety, and molecular pharmacology: potential for collaboration

TO THE EDITOR: In an effort to enhance our ability to detect novel adverse events in a timely manner, we are studying computational signal detection algorithms, also known as data mining algorithms (DMAs), in large postmarketing safety databases to determine whether these computational methods might usefully supplement our traditional surveillance strategies. We are in the process of publishing our findings.^{1,2} DMAs include simple disproportionality analysis (eg, proportional reporting ratios [PRRs]) and algorithms that use additional statistical adjustments and Bayesian modeling (eg, multi-item gamma-Poisson shrinker [MGPS]). These DMAs are being applied to spontaneous reporting system databases to screen for disproportionate statistical dependencies between drugs and events. If there is sufficient correlation between the aforementioned statistical dependencies (potential signals) and causal drug–event associations, these techniques may enhance our ability to detect adverse events caused by medicines. We propose a possible extended application of these techniques.

A structure–adverse event relationship, not previously investigated, was identified from laboratory research and theoretical modeling performed in Japan during the years 1998–2000.^{3–5} The authors noted from their *in vitro* and *in vivo* research that the diethyl amino methyl moiety contained in drugs from disparate pharmacologic/therapeutic classes marketed in Japan (other than antipsychotics) seemed to play a key role in the induction of parkinsonism and/or catalepsy. In their publications, they supported their findings with published anecdotal case reports. We questioned whether application of DMAs to a large postmarketing safety database could be of potential use in assisting researchers to identify drugs with relevant pharmacophores (eg, molecular structural features) for structure–adverse effect modeling. MGPS and PRRs were used to data mine on drug–event combinations (DECs) in the Food and Drug Administration Adverse Events Reporting System database relating to an association of parkinsonism with drugs containing the diethyl amino methyl moiety. For 9 of the 12 drugs listed in the research publications that were not antipsychotics (ie, tiapride, haloperidol),^{3–5} there were reports of parkinsonism-related events in the database. Antipsychotics were not evaluated because this class of drugs is known to be associated with parkinsonism. For 3 drugs, there were reports but no signals of disproportionate reporting (SDRs). Table 1 demonstrates the number of reports and the associated level of disproportionate reporting. An SDR for a DEC means that the reporting frequency of that DEC was higher than might be expected if drug and event were independently distributed in the database and reflects reporting activity and not necessarily causality. Six DEC generated an SDR with PRRs and 4 of the 6 (amoxapine, cinnarizine, domperidone, metoclopramide) signaled with PRRs and/or MGPS in advance of the publication dates of the papers. None signaled with MGPS alone.

It appears that there is a potential for collaboration between drug safety experts and molecular pharmacologists using DMAs as tools for the

Table 1. Number of Reports and Associated Level of Disproportionate Reporting by Drug and Event^a

Drug	Class	Year of First Report	Year of First SDR ^b	
			PRR ^c	MGPS
Amiodarone	antiarrhythmic	1990	parkinsonism 1998 (n = 2)	NS
Amoxapine	antidepressant	1981	dystonia 1981 (n = 1) extrapyramidal disorder 1982 (n = 9)	extrapyramidal disorder 1983 (n = 17) dystonia 1984 (n = 12)
Cinnarizine	cerebrovascular disorders	1993	extrapyramidal disorder 1993 (n = 1)	NS
Domperidone	antiemetic	1996	extrapyramidal disorder 1996 (n = 1) Parkinson's disease NOS 1998 (n = 1)	Parkinson's disease NOS 2002 (n = 4)
Flunarizine	cerebrovascular disorders	2001	extrapyramidal disorder 2001 (n = 1)	NS
Metoclopramide	antiemetic	1977	extrapyramidal disorder 1977 (n = 1) dystonia 1981 (n = 2)	dystonia 1982 (n = 6) extrapyramidal disorder 1982 (n = 9)

MGPS = multi-item gamma-Poisson shrinker; NOS = not otherwise specified; NS = no SDR between drug and event; PRR = proportional reporting ratio; SDR = signal of disproportionate reporting.
^aSDR persisted over time and/or with accumulation of additional cases.
^bFor event(s) signaled by either algorithm (n cases).
^cFor purposes of detecting SDRs in pharmacovigilance, a case count threshold is frequently used (eg, n > 2).

identification of drugs from diverse pharmacologic/therapeutic classes that may contain pharmacophores relevant to the analysis of structure–adverse effect relationship. In practice, a set of drugs signaling for an adverse event could be the basis for initial screening for possible pharmacophore modeling and structure–activity screens to identify patterns that might be fruitful areas for further scientific research. We encourage further investigation in this area.

Manfred Hauben MD MPH

Medical Director, Risk Management Strategy
Pfizer Inc.

New York, New York

Department of Medicine

New York University School of Medicine

New York, New York

Departments of Pharmacology and Community and Preventive Medicine

New York Medical College

Valhalla, New York

Lester Reich MD

Medical Director, Risk Management Strategy
Pfizer Inc.

235 East 42nd Street

New York, New York 10017-5755

fax 212/808-8679

lester.reich@pfizer.com

Published Online, 9 Nov 2004, www.theannals.com
DOI 10.1345/aph.1E373

REFERENCES

1. Hauben M, Reich L. Safety related drug-labeling changes: findings from two data mining algorithms. *Drug Saf* 2004;27:735-44.
2. Hauben M, Reich L. Drug-induced pancreatitis: lessons in data mining. *Br J Clin Pharmacol* 2004;58:560-2.
DOI 10.1111/j.1365-2125.2004.02176
3. Matsui A, Matsuo H, Hitomi T, Sasaki S, Maeda M, Sawada Y. Prediction of catalepsies induced by amiodarone, aprindine and procaine: similarity in conformation of diethylaminoethyl side chain. *J Pharmacol Exp Ther* 1998;287:725-32.
4. Matsuo H, Matsui A, Risa N, Takanaga H, Inoue N, Hattori F, et al. Propiverine-induced parkinsonism: a case report and a pharmacokinetic/pharmacodynamic study in mice. *Pharm Res* 2000;17:565-71.
5. Sawada Y, Nasu R, Matsuo H, Ohtani H. Relationship between structure and drug-induced parkinsonism (letter). *Ann Pharmacother* 2000;34:668.
DOI 10.1345/aph.19253

Anaphylaxis to intravenous sinistrin

TO THE EDITOR: Inulin/sinistrin clearance is the gold standard for the determination of glomerular filtration rate (GFR). Sinistrin, an analog of inulin, has better water solubility. Anaphylaxis to intravenous administration of inulin/sinistrin is rarely reported. We report an anaphylactic reaction in a healthy male who received sinistrin.

Case Report. A healthy, 25-year-old male Chinese volunteer participated in a Phase I clinical trial with assessment of GFR by sinistrin (Inutest, Fresenius Pharma Austria, Graz). No atopy or allergies were known from his history. No investigations with inulin/sinistrin had been performed previously. He was not taking any medication and had not received any investigational medication prior to sinistrin.

After an overnight fast, sinistrin 2 g in 45 mL of NaCl 0.9% solution with 100 mg of sodium lactate was administered intravenously. After 4 minutes, dyspnea without stridor and a prickling and itching on neck and face appeared. Angioedema of the lips and symmetrical urticarial erythemas on the patient's head, neck, trunk, and upper extremities were observed.

The infusion of sinistrin was stopped, and oxygen (3 L/min) was administered. Prednisolone 100 mg and dimethindene 4 mg (antihistamine agent) were administered intravenously. An electrocardiogram (ECG) indicated an incomplete right bundle branch block with negative T waves ($V_{1,2}$). U waves appeared in leads $V_{3,5}$ (Figure 1). Hypokalemia (3.1 mEq/L) was measured. After one hour, symptoms subsided and the ECG normalized. After 24 days, skin prick testing for sinistrin was immediately positive (1:1 solution).

Discussion. Our patient's symptoms (generalized urticaria, angioedema, dyspnea) and a positive skin prick test indicated an anaphylactic reaction to intravenous sinistrin. Use of the Naranjo probability scale indi-

cated a probable relationship between anaphylaxis and sinistrin.¹ Bronchospasm and, thus, pulmonary hypertension, leading to acute right ventricular stress, may have caused the incomplete right bundle branch block. Hypokalemia and U waves in the ECG were likely to be due to endogenous epinephrine.

To our knowledge, only one other case of anaphylaxis upon intravenous administration of sinistrin has been reported.² A 45-year-old white patient with Wegener's granulomatosis developed anaphylaxis and cardiorespiratory arrest after administration of bolus intravenous sinistrin (2.5 g in 10 mL of solution). A letter has reported anaphylaxis occurring on oral ingestion of inulin in vegetables and processed foods.³

Other polycarbohydrates used in medicine, such as the plasma expanders dextran, gelatin, and hydroxyethyl starch (HES), have low incidences of anaphylaxis (0.273%, 0.345%, and 0.058%, respectively).^{4,5} Dextran anaphylaxis is caused by circulating dextran-reactive immunoglobulin G found in most adults.⁴ Gelatin may lead to histamine release from mast cells.⁴ The mechanism of HES anaphylaxis is unclear; presumably HES-reactive antibodies preformed in humans do not exist or are extremely rare.^{4,6} There are no known cross-reactivities between these substances and inulin/sinistrin.⁵

Anaphylaxis to intravenous sinistrin in healthy individuals, as in our case, seems to be very rare or underreported. The underlying mechanism of inulin/sinistrin anaphylaxis is still unclear. Skin prick tests suggest that the immediate-type hypersensitivity reaction may be immunoglobulin E-mediated.

Richard Fux MD

Clinical Pharmacologist

Abteilung Klinische Pharmakologie

Institut für Pharmakologie und Toxikologie

Universitätsklinikum Tübingen

Tübingen, Germany

Figure 1. ECG during (upper panel) and after (lower panel) clinical symptoms: incomplete right bundle branch block with negative T waves (V_1 and V_2), U waves in leads $V_{3,5}$.

Dermatologist
Universitäts-Hautklinik Tübingen
Universitätsklinikum Tübingen

Tilo Biedermann MD

Dermatologist
Universitäts-Hautklinik Tübingen
Universitätsklinikum Tübingen

Tine Sander-Wiecker MD

Clinical Pharmacologist
Abteilung Klinische Pharmakologie
Institut für Pharmakologie und Toxikologie
Universitätsklinikum Tübingen

Klaus Mörke MD

Clinical Pharmacologist, Head
Abteilung Klinische Pharmakologie
Institut für Pharmakologie und Toxikologie
Universitätsklinikum Tübingen
Koordinierungszentrum Klinische Studien am Universitätsklinikum
Tübingen (KKS-UKT gGmbH)
Otfried-Müller-Str. 45, D-72076 Tübingen, Germany
fax 49 (0)7071 29-5035
christoph.gleiter@med.uni-tuebingen.de

Christoph H Gleiter MD

Published Online, 9 Nov 2004, www.theannals.com
DOI 10.1345/aph.1E267

REFERENCES

1. Naranjo CA, Busto U, Sellers EM, Sandor P, Ruiz I, Roberts EA, et al. A method for estimating the probability of adverse drug reactions. *Clin Pharmacol Ther* 1981;30:239-45.
2. Chandra R, Barron JL. Anaphylactic reaction to intravenous sinistrin (Inu-test). *Ann Clin Biochem* 2002;39(pt 1):76.
3. Gay-Crosier F, Schreiber G, Hauser C. Anaphylaxis from inulin in vegetables and processed foods (letter). *N Engl J Med* 2000;342:1372.
4. Dieterich HJ, Kraft D, Sirtl C, Laubenthal H, Schimetta W, Polz W, et al. Hydroxyethyl starch antibodies in humans: incidence and clinical relevance. *Anesth Analg* 1998;86:1123-6.
5. Lehmann G, Asskali F, Förster H. Severe adverse event following iv administration of 10 ml 6% Dextran60 (0.6 g) in a healthy volunteer. *Anaesthesist* 2002;51:820-4.
6. Kraft D, Sirtl C, Laubenthal K, Scheiner O, Parth E, Dieterich HJ, et al. No evidence for the existence of preformed antibodies against hydroxyethyl starch in man. *Eur Surg Res* 1992;24:138-42.

Comment: renal protective effects of angiotensin receptor blockers

TO THE EDITOR: Coyle et al.¹ recently prepared an overview of the evidence supporting the nephroprotective effects of angiotensin II receptor blockers (ARBs). They reviewed several large trials comparing ARBs with placebo or active control.^{2,3} Based on these data, ARBs unarguably slow the progression of diabetic nephropathy in patients with and without chronic renal insufficiency (CRI) as evidenced by a reduction in the composite outcome of doubling of the serum creatinine level, development of end-stage renal disease (ESRD), and death. Three matters are worthy of mention with respect to the review. First, patients with diabetic nephropathy are at extremely high risk for future cardiovascular events, including myocardial infarction and cardiovascular death. In fact, cardiovascular events occurred more often than renal outcomes in these trials. ARBs have not been shown to reduce either of these outcomes in a wide range of patients. On the other hand, angiotensin-converting enzyme (ACE) inhibitors are both nephroprotective and cardioprotective, decreasing all major cardiovascular endpoints in high-risk patients.

Comments on articles previously published are submitted to the authors of those articles. When no reply is published, either the author chose not to respond or did not do so in a timely fashion. Comments and replies are not peer reviewed.—ED.

Second, evidence emerging after the publication of the 2 largest renal protection trials questions whether ARBs will decrease ESRD in clinical practice.^{2,3} The RENAAL (Reduction of Endpoints in NIDDM with the Angiotensin II Antagonist Losartan) study was conducted at 250 centers in 28 countries in Asia, Europe, Central America, South America, and North America. At the end of follow-up, the primary endpoint was reduced by 16%. This reduction was driven largely by a 35% reduction in the Asian population, with smaller, nonsignificant reductions on all other continents. In the IDNT (Irbesartan Diabetic Nephropathy Trial), there was a 20% reduction in the primary endpoint. However, there was no significant difference in progression to ESRD or death, and patients receiving irbesartan suffered a similar number of cardiovascular events as patients who received placebo. The major benefit of irbesartan was its effect on a surrogate endpoint, which was the change in serum creatinine.

Lastly, Figure 1 in the review presents a proposed treatment algorithm for patients with diabetes and microalbuminuria or nephropathy. Step 1 in the algorithm suggests that glycemic control should be addressed first. Blood pressure is considered only after glycemic control has been achieved. This is of concern given the findings from the UKPDS 38 (United Kingdom Prospective Diabetes Study)⁴ and, more recently, the VALUE (Valsartan Antihypertensive Long-term Use Evaluation) trial.⁵ In the UKPDS, aggressive blood pressure control was associated with reductions in mortality, all diabetes-related events, microvascular disease, and stroke of 18%, 24%, 37%, and 44%, respectively. Conversely, aggressive control of glycemia reduced all diabetes-related events by 12% and microvascular disease by 19%, but did not affect macrovascular disease or death. Furthermore, in the VALUE trial, compared with patients whose blood pressure was lowered to <140/90 mm Hg by 6 months, patients with uncontrolled blood pressure experienced a 21% increase in mortality, 27% increase in cardiovascular events, and 50% increase in cerebrovascular accidents. Consequently, hypertension should receive the bulk of attention since it is expected to have a more profound impact on both micro- and macrovascular disease and because even a minimal delay in adequate control could be detrimental. Blood pressure should be lowered aggressively to <130/80 mm Hg with agents that have been shown to prevent cardiovascular events in this population, namely ACE inhibitors.

Benjamin J Epstein PharmD

Postdoctoral Fellow
Departments of Pharmacy Practice and Family Medicine
Colleges of Pharmacy and Medicine
University of Florida
Gainesville, Florida 32601-6430
fax 352/392-7766
epstein@chfm.ufl.edu

Published Online, 9 Nov 2004, www.theannals.com
DOI 10.1345/aph.1E182a

REFERENCES

1. Coyle JD, Gardner SF, White MW. The renal protective effects of angiotensin II receptor blockers in type 2 diabetes mellitus. *Ann Pharmacother* 2004;38:1731-8. DOI 10.1345/aph.1E182
2. Brenner BM, Cooper ME, de Zeeuw D, Keane WF, Mitch WE, Parving HH, et al. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001;345:861-9.
3. Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, et al., for the Collaborative Study Group. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001;34:851-60.
4. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group. *BMJ* 1998;317:703-13.
5. Julius S, Kjeldsen SE, Weber M, Brunner HR, Ekman S, Hansson L, et al., for the VALUE trial group. Outcomes in hypertensive patients at high cardiovascular risk treated with regimens based on valsartan or amlodipine: the VALUE randomised trial. *Lancet* 2004;363:2022-31.

AUTHORS' REPLY: We thank Dr. Epstein for his comments in response to our review of the renal protective effects of ARBs in patients with type 2 diabetes.¹ We would like to address each of his 3 major points.

First, Dr. Epstein suggested that ACE inhibitors are clearly preferred over ARBs for cardiovascular risk reduction in patients with diabetic nephropathy. This position is becoming increasingly difficult to defend. There is a significant, growing body of literature showing that ARBs also decrease cardiovascular risk, that the magnitude of their overall effect is similar to that of ACE inhibitors, and that they have a more favorable adverse effect profile than ACE inhibitors.²⁻¹⁰ For example, VALIANT recently provided a head-to-head comparison of the effects of ARB monotherapy (losartan), ACE inhibitor monotherapy (captopril), and combination ARB and ACE inhibitor therapy (valsartan plus captopril) on mortality in a high-risk patient population with acute myocardial infarction complicated by heart failure, left ventricular systolic dysfunction, or both.¹⁰ All-cause mortality, cardiovascular mortality, recurrent myocardial infarction, and hospitalizations for heart failure were similar in all treatment groups, while the lowest incidence of adverse events occurred in the losartan monotherapy treatment group.

Other recent clinical trials also support ARB use to reduce cardiovascular risk without providing a direct comparison with ACE inhibitors.^{11,12} In our opinion, available data support the use of either ARBs or ACE inhibitors to provide renal and cardiovascular protection in patients with type 2 diabetes as indicated in the treatment algorithm in our manuscript.

Dr. Epstein's second point primarily questions the ability of ARBs to decrease ESRD in clinical practice. Many aspects of his argument regarding specific aspects of individual studies are correct. However, in our opinion, the entire body of evidence regarding the renal protective effects of ARBs and ACE inhibitors strongly suggests that both classes of agents are renoprotective. We refer readers to the *K/DOQI Clinical Practice Guidelines on Hypertension and Antihypertensive Agents in Chronic Kidney Disease* for a comprehensive summary of current data on this topic.¹³ The American Diabetes Association's 2004 Clinical Practice Recommendations are also consistent with our view.¹⁴

Dr. Epstein's third point questions the appropriateness of good glycemic control as the first step in our treatment algorithm. We agree that strict blood pressure control is more important than strict glycemic control in protecting the kidneys of patients with diabetes. What we intended to convey in our algorithm is that achieving strict glycemic control contributes to the lowering of blood pressure and to renal protection in patients with diabetes. Simultaneously achieving glycemic and blood pressure control should be advocated.

Our review article was narrowly focused on the renal protective effects of ARBs in patients with type 2 diabetes since it was intended to serve as an update to the excellent review of the renoprotective effects of ACE inhibitors (and other drugs) that appeared previously in *The Annals*.¹⁵ We hope that Dr. Epstein's comments and our reply have helped clarify the current status of some potentially controversial issues related to this topic.

James D Coyle PharmD

Assistant Professor of Clinical Pharmacy
Division of Pharmacy Practice and Administration
College of Pharmacy
The Ohio State University
Columbus, Ohio

Stephanie F Gardner PharmD EdD

Professor and Dean
College of Pharmacy
UAMS
Little Rock, Arkansas

C Michael White PharmD

Associate Professor of Pharmacy
School of Pharmacy
University of Connecticut
80 Seymour Street
Hartford, Connecticut 06102-5037
fax 860/545-2277
cmwhite@harthosp.org

Published Online, 9 Nov 2004, www.theannals.com
DOI 10.1345/aph.1E182b

REFERENCES

1. Coyle JD, Gardner SF, White CM. The renal protective effects of angiotensin II receptor blockers in type 2 diabetes mellitus. *Ann Pharmacother* 2004;38:1731-8. DOI 10.1345/aph.1E182
2. Pitt B, Segal R, Martinez FA, Meurers G, Cowley AJ, Thomas I, et al. Randomised trial of losartan versus captopril in patients over 65 with heart failure (Evaluation of Losartan in the Elderly Study, ELITE). *Lancet* 1997;349:747-52.
3. Pitt B, Poole-Wilson PA, Segal R, Martinez FA, Dickstein K, Camm AJ, et al. Effect of losartan compared with captopril on mortality in patients with symptomatic heart failure: randomized trial—the Losartan Heart Failure Survival Study, ELITE II. *Lancet* 2000;355:1582-7.
4. Cohn JN, Tognoni G. A randomized trial of the angiotensin-receptor blocker valsartan in chronic heart failure. *N Engl J Med* 2001;345:1667-75.
5. Maggioni AP, Anand I, Gottlieb SO, Latini R, Tognoni G, Cohn JN. Effects of valsartan on morbidity and mortality in patients with heart failure not receiving angiotensin-converting enzyme inhibitors. *J Am Coll Cardiol* 2002;40:1414-21.
6. Pfeffer MA, Swedberg K, Granger CB, Held P, McMurray JJ, Michelson EL, et al. Effects of candesartan on mortality and morbidity in patients with chronic heart failure: the CHARM-Overall programme. 2003;362:759-66.
7. McMurray JJ, Ostergren J, Swedberg K, Granger CB, Held P, Michelson EL, et al. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function taking angiotensin-converting-enzyme inhibitors: the CHARM-Added trial. *Lancet* 2003;362:767-71.
8. Granger CB, McMurray JJ, Yusuf S, Held P, Michelson EL, Olofsson B, et al. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function intolerant to angiotensin-converting-enzyme inhibitors: the CHARM-Alternative trial. *Lancet* 2003;362:772-6.
9. Yusuf S, Pfeffer MA, Swedberg K, Granger CB, Held P, McMurray JJ, et al. Effects of candesartan in patients with chronic heart failure and preserved left-ventricular ejection fraction: the CHARM-Preserved trial. *Lancet* 2003;362:777-81.
10. Pfeffer MA, McMurray JJ, Velazquez EJ, Rouleau JL, Kober L, Maggioni AP, et al. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med* 2003;349:1893-906.
11. Dahlöf B, Devereux RB, Kjeldsen SE, Julius S, Beevers G, de Faire U, et al. Cardiovascular morbidity and mortality in the losartan intervention for endpoint reduction in hypertension study (LIFE): a randomized trial against atenolol. *Lancet* 2002;359:995-1003.
12. Papademetriou V, Varsang C, Elmfeldt D, Hofman A, Lithell H, Olofsson B, et al. Stroke prevention with the angiotensin II type 1-receptor blocker candesartan in elderly patients with isolated systolic hypertension: the study on cognition and prognosis in the elderly (SCOPE). *J Am Coll Cardiol* 2004;44:1175-80.
13. National Kidney Foundation. K/DOQI clinical practice guidelines on hypertension and antihypertensive agents in chronic kidney disease. *Am J Kidney Dis* 2004;43(suppl 1):S1-290.
14. American Diabetes Association 2004 Clinical Practice Recommendations. *Diabetes Care* 2004;27(suppl 1):S1-150.
15. Vivian EM, Goebig ML. Slowing the progression of renal disease in diabetic patients. *Ann Pharmacother* 2001;35:452-63. DOI 10.1345/aph.10067

Letters are subject to review prior to acceptance. They should address areas related to pharmacy practice, research, or education, or articles recently published. Corrections of previously published material also are accepted. Letters are limited to no more than five authors. In cases where adverse drug effects are described, the Naranjo ADR probability scale should be used to determine the likelihood that the adverse effect was drug-related (*Clin Pharmacol Ther* 1981;30:239-45). Text: limit 500 words. References: limit 5. Art: limit 1 table or figure.